Dimensions closed: 8.5" x 11" Spread: 17" x 11"



Billing and Coding Considerations for BLINCYTO®

This Information Sheet is intended to help healthcare professionals understand the key billing and coding considerations for BLINCYTO[®] and its related services and supplies when using the Food and Drug Administration (FDA)-approved dosing options across treatment settings.

Updates regarding Medicare Home Infusion Therapy Benefit:

- **1.** Starting January 1, 2021, Medicare implemented the permanent home infusion therapy benefit that provides separate Part B coverage and payment for qualified home infusion therapy services¹
 - · Medicare updated the codes used to report the provision of home infusion therapy services
 - The new codes differentiate new visits vs subsequent visits for home infusion therapy services
 - Claims for home infusion therapy services will be billed separately from the drug, pump, and other supplies. These services must be reported to the A/B Medicare Administrative Contractor (MAC), and are reimbursed by Medicare at rates set by the Medicare Physician Fee Schedule. Claims for the drug, pump, and supplies should continue being sent to the Durable Medical Equipment (DME) MAC and are payable under the Prosthetics, Orthotics, and Supplies (DMEPOS) Fee Schedule^{1,2}
 - Home infusion therapy services are equal to 5 hours per calendar day, billed in 15 minute increments

Please see pages 10 and 11 for sample claim forms showing coding changes that may be appropriate to report services for Medicare beneficiaries receiving BLINCYTO® treatment via home infusion

2. Due to COVID-19 Public Health Emergency (PHE), Medicare temporarily revised the definition of direct supervision to include the virtual presence of the supervising physician or other qualified healthcare provider using real-time, interactive audio and video telecommunications technology through to December 31, 2024³

Medicare sequestration has been fully reinstated beginning with the third quarter of 2022 and as such, the Medicare portion of payment rates are reduced by 2%.4

Please note that the information in this resource is intended to be educational and is not a guarantee of reimbursement. Coverage, coding, and billing requirements vary by health plan so be sure to check with individual payers for detailed guidance.

INDICATIONS

BLINCYTO® (blinatumomab) is indicated for the treatment of CD19-positive B-cell precursor acute lymphoblastic leukemia (ALL) in adult and pediatric patients one month and older with:

- Philadelphia chromosome-negative disease in the consolidation phase of multiphase chemotherapy
- Minimal residual disease (MRD) greater than or equal to 0.1% in first or second complete remission
- Relapsed or refractory disease

IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGICAL TOXICITIES including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

- Cytokine Release Syndrome (CRS), which may be life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® and treat with corticosteroids as recommended.
- Neurological toxicities, including immune effector cell-associated neurotoxicity syndrome (ICANS) which may be severe, life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® as recommended.

Please see accompanying BLINCYTO® full Prescribing Information, including **BOXED WARNINGS**. Please see additional Important Safety Information on pages 18-19.

BLINCYTO®

Billing Information Sheet

Hospital Inpatient (HIP) Site of Service - Multiple Payers (Medicare and Non-Medicare)

Item	Revenue Code ^{5,6,*}	Coding Information (ICD-10-CM ⁷ /HCPCS ⁸ /CPT ⁹ /ICD-10-PCS ¹⁰)	Notes
Diagnosis: Encounter for drug therapy and ALL	N/A	Z51.12 Encounter for antineoplastic immunotherapy AND C91.00 Acute lymphoblastic leukemia not having achieved remission/failed remission OR C91.01 Acute lymphoblastic leukemia, in remission OR C91.02 Acute lymphoblastic leukemia, in relapse	Report the appropriate ICD-10-CM diagnosis code(s) to describe the patient's condition.
Drug: BLINCYTO® and external infusion pump (EIP)	Report the appropriate revenue code for the cost center in which the service is performed; eg, • Medicare: 0250 General pharmacy • Other payers: 0250 or 0636 Drugs requiring detailed coding (if required by a given payer)	J9039 Injection, blinatumomab, 1 mcg	
	Report the appropriate revenue code for the cost center in which the service is performed; eg, • 0290 DME	E0791 Parenteral infusion pump, stationary, single or multi-channel E0776 IV pole	
Administration: Continuous intravenous infusion (CIVI) via EIP	Report the appropriate revenue code for the cost center in which the service is performed; eg, • 0261 IV therapy: Infusion pump	3E03305 Introduction of other antineoplastic into peripheral vein, percutaneous approach† OR 3E04305 Introduction of other antineoplastic into central vein, percutaneous approach† 96416 Chemotherapy administration, IV infusion technique; initiation of prolonged chemotherapy infusion (more than 8 hours) requiring use of a portable or implantable pump OR 96521 Refilling and maintenance of a portable pump	

Coding Information Definitions:

ICD-10-CM – International Classification of Diseases, 10th Revision, Clinical Modification

HCPCS - Healthcare Common Procedure Coding System

CPT - Current Procedural Terminology


ICD-10-PCS – International Classification of Diseases, 10th Revision, Procedure Coding System

*This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.

†The previous ICD-10-PCS codes that described the administration of BLINCYTO® (XW03351 and XW04351) have been deleted and should not be used for dates of service on or after October 1, 2021.



Sample UB-04 (CMS-1450) Form: Hospital Inpatient Administration



^{*}This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.

This sample form is intended as a reference for coding and billing for product and associated services. It is not intended to be directive; the use of the recommended codes does not guarantee reimbursement. Healthcare providers may deem other codes or policies more appropriate and should select the coding options that most accurately reflect their internal system guidelines, payer requirements, practice patterns, and the services rendered. Healthcare providers are responsible for ensuring the accuracy and validity of all billing and claims for appropriate reimbursement.

[†]The previous ICD-10-PCS codes that described the administration of BLINCYTO® (XW03351 and XW04351) have been deleted and should not be used for dates of service on or after October 1, 2021.

Hospital Outpatient Department (HOPD) - Multiple Payers (Medicare and Non-Medicare)

Item	Revenue Code ^{5,6,*}	Coding Information (ICD-10-CM ⁷ /CPT ⁹ /HCPCS ⁸ /NDC ¹¹)	Notes
Diagnosis: Encounter for drug therapy and ALL	N/A	Z51.12 Encounter for antineoplastic immunotherapy AND C91.00 Acute lymphoblastic leukemia not having achieved remission OR C91.01 Acute lymphoblastic leukemia, in remission OR C91.02 Acute lymphoblastic leukemia, in relapse	Report the appropriate ICD-10-CM diagnosis code(s) to describe the patient's condition.
Procedure: Administration via CIVI using an EIP	Report the appropriate revenue code for the cost center in which the service is performed; eg, • 0261 IV therapy: Infusion pump • 026x IV therapy	96416 Chemotherapy administration, IV infusion technique; initiation of prolonged chemotherapy infusion (more than 8 hours) requiring use of a portable or implantable pump 0R 96521 Refilling and maintenance of portable pump 0R G0498 Chemotherapy administration, IV infusion technique; initiation of infusion in the office/clinic setting using office/clinic pump/ supplies, with continuation of the infusion in the community setting (eg, home, domiciliary, rest home, or assisted living) using a portable pump provided by the office/clinic; includes follow-up office/clinic visit at the conclusion of the infusion	CPT codes may be used to report the CIVI procedures associated with BLINCYTO® to the Part A/B MAC and non-Medicare payers. For Medicare patients, HCPCS code G0498 will replace CPT and HCPCS codes (96416, E0781, and 99211–99215) previously used to bill for prolonged infusion services when the CIVI is started in the HOPD. It does not apply to BLINCYTO® when the CIVI is started in the inpatient setting or via home infusion. 8,9,12 Certain payers may not recognize G0498 and require itemization of specific items, instead. The healthcare provider should consult the payer or MAC to determine which code is most appropriate for administration of BLINCYTO®. If the clinic bills the G-code to the MAC, the cost of the pump and supplies is bundled and should not be billed separately to the DME MAC. 13
Drug: BLINCYTO®	Report the appropriate revenue code for the cost center in which the service is performed; eg, • Medicare: 0636 Drug requiring detailed coding • Other payers: 0250 or 0636 General pharmacy (if required by a given payer)	J9039 Injection, blinatumomab, 1 mcg JW Discarded drug/not administered to any patient JZ Zero drug amount discarded/not administered to any patient JG Drug or biological acquired with 340B Drug Pricing Program discount TB Drug or biological acquired with 340B Drug Pricing Program discount	Medicare policies reflect the code for BLINCYTO® (J9039 per 1 mcg) and has a maximum utilization of 210 units per date of service (based on prescribing information). However, coding and coverage requirements may vary by payer. Like many payers, Medicare requires the use of the modifier JW and JZ, which provides payment for the amount of drug or biologic discarded, as well as for the dose administered, up to the amount of the drug or biologic as indicated on the vial or label for a single-dose vial (SDV). Note: Effective for dates of service on or after July 1, 2023, Medicare Part B claims require the use of the new JZ modifier for single-use vials and containers when there are no discarded drug amounts. Medicare claims also continue to require the use of the JW modifier (Drug amount discarded/not administered to any patient) for drugs and biologicals that are separately payable under Medicare Part B with discarded amounts from single-dose containers. Starting January 1, 2024, CMS is requiring all 340B covered entities, including hospital-based and nonhospital-based entities, that submit claims for separately payable Part B drugs and biologicals to report modifier "JG" or "TB" on claim lines for drugs acquired through the 340B Drug Pricing Program. Starting January 1, 2025, 340B covered entities must report the "TB" modifier on claims. Calmin.
	N/A	NDC: 55513016001 BLINCYTO® 35 mcg lyophilized powder, SDV IV solution stabilizer, 10 mL SDV	Some payers (eg, Medicaid) may require listing the NDC in addition to the HCPCS J-code. When reporting the NDC on claims, use the 11-digit NDC in the 5-4-2 format. Insert a leading zero in the appropriate section to complete the 5-4-2 digit format. Remove the dashes prior to entering the NDC on the claim form.

Coding Information Definition: NDC – National Drug Code



Hospital Outpatient Department (HOPD) - Multiple Payers (Medicare and Non-Medicare) (continued)

Item	Revenue Code ^{5,6,*}	Coding Information (HCPCS ⁸)	Notes
DME: EIP and supplies	Report the appropriate revenue code for the cost center in which the service is performed; eg, 0290 DME	E0779 Ambulatory infusion pump, mechanical, reusable, for infusion 8 hours or greater OR E0781 Ambulatory infusion pump, single or multiple channels, electric or battery operated, with administrative equipment, worn by patient OR A4222 Infusion supplies for external drug infusion pump, per cassette or bag Modifiers for use with E-codes for IV pump -KD Drug or biologic infused through DME -RR Rental -KH DMEPOS item, initial claim, purchase or first rental month -KI DMEPOS item, second or third rental months -KJ DMEPOS item, parenteral enteral nutrition (pen) pump or capped rental, fourth to 15th rental months	Please note that Medicare specifically requires DMEPOS accreditation in order to bill a DME MAC. Non-Medicare payers may allow billing for all services and supplies under a medical or other benefit. Report the appropriate EIP code and appropriate modifier(s) as documented in the medical record. Modifiers may be used to provide additional detail when billing for the EIP to the DME MAC. ⁸ Note: Drug administration codes may get billed to the MAC and the E-codes may get billed separately to the DME MAC. Report any supplies as necessary.

^{*}This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.


^{*}This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.

†Reporting policies for discarded units for payers other than traditional fee-for-service Medicare may vary; providers should check with their specific plans about policies related to billing for discarded drug and use of the JW and JZ modifiers.

BLINCYTO®

Billing Information Sheet

Sample UB-04 (CMS-1450) Form: Hospital Outpatient Administration



*This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.

This sample form is intended as a reference for coding and billing for product and associated services. It is not intended to be directive; the use of the recommended codes does not guarantee reimbursement. Healthcare providers may deem other codes or policies more appropriate and should select the coding options that most accurately reflect their internal system guidelines, payer requirements, practice patterns, and the services rendered. Healthcare providers are responsible for ensuring the accuracy and validity of all billing and claims for appropriate reimbursement.



Physician Office - Multiple Payers (Medicare and Non-Medicare)

Item	Coding Information (ICD-10-CM ⁷ /CPT ⁹ /HCPCS ⁸ /NDC ¹¹)	Notes
Diagnosis: Encounter for drug therapy and ALL	Z51.12 Encounter for antineoplastic immunotherapy AND C91.00 Acute lymphoblastic leukemia not having achieved remission OR C91.01 Acute lymphoblastic leukemia, in remission OR C91.02 Acute lymphoblastic leukemia, in relapse	Report the appropriate ICD-10-CM code(s) to describe the patient's condition.
Procedure: Administration via CIVI using an EIP	96416 Chemotherapy administration, intravenous infusion technique; initiation of prolonged chemotherapy infusion (more than 8 hours) requiring use of a portable or implantable pump OR 96521 Refilling and maintenance of portable pump OR G0498 Chemotherapy administration, IV infusion technique; initiation of infusion in the office/clinic setting using office/clinic pump/supplies, with continuation of the infusion in the community setting (eg, home, domiciliary, rest home, or assisted living) using a portable pump provided by the office/clinic; includes follow-up office/clinic visit at the conclusion of the infusion	CPT codes may be used to report the CIVI procedures associated with BLINCYTO® to the Part A/B MAC and non-Medicare payers. For Medicare patients, HCPCS code G0498 will replace CPT codes and HCPCS (96416, E0781, and 99211–99215) previously used to bill for prolonged infusion services when the CIVI is started in the physician office. It does not apply to BLINCYTO® when the CIVI is started in the inpatient setting or via home infusion. 8,912 Some payers may not recognize G0498 and require itemization of specific items, instead. The healthcare provider should consult the payer or MAC to determine which code is most appropriate for administration of BLINCYTO®.
Drug: BLINCYTO®	J9039 Injection, blinatumomab, 1 mcg JW Discarded drug/not administered to any patient JZ Zero drug amount discarded/not administered to any patient	Medicare requires use of the HCPCS code in the physician office setting ¹⁸ and has a maximum utilization of 210 units per date of service (based on prescribing information). ¹⁹ However, coding requirements may vary by payer. Like many payers, Medicare requires the use of the modifier JW and JZ, which provides payment for the amount of drug or biologic discarded, as well as for the dose administered, up to the amount of the drug or biologic as indicated on the vial or label for an SDV. ¹⁵ Note: Effective for dates of service on or after July 1, 2023, Medicare Part B claims require the use of the new JZ modifier for single-use vials and containers when there are no discarded drug amounts. Medicare claims also continue to require the use of the JW modifier (Drug amount discarded/not administered to any patient) for drugs and biologicals that are separately payable under Medicare Part B with discarded amounts from single-dose containers.*
	NDC: 55513016001 BLINCYTO® 35 mcg lyophilized powder, SDV IV solution stabilizer, 10 mL SDV	Some payers (eg, Medicaid) may require listing the NDC in addition to the HCPCS J-code. When reporting the NDC on claims, use the 11-digit NDC in the 5-4-2 format. ¹⁷ Insert a leading zero in the appropriate section to complete the 5-4-2 digit format. Remove the dashes prior to entering the NDC on the claim form.
DME: EIP and supplies	E0779 Ambulatory infusion pump, mechanical, reusable, for infusion 8 hours or greater ²² E0781 Ambulatory infusion pump, single or multiple channels, electric or battery operated, with administrative equipment, worn by patient ²² G0498 Chemotherapy administration, IV infusion technique; initiation of infusion in the office/clinic setting using office/clinic pump/supplies, with continuation of the infusion in the community setting (eg, home, domiciliary, rest home, or assisted living) using a portable pump provided by the office/clinic; includes follow-up office/clinic visit at the conclusion of the infusion A4222 Infusion supplies for external drug infusion pump, per cassette or bag Modifiers for EIP -KD Drug or biologic infused through DME -RR Rental -KH DMEPOS item, initial claim or first rental month -KI DMEPOS item, second or third rental months -KJ DMEPOS item, fourth to 15th rental months	Report the appropriate EIP code and appropriate modifier(s) as documented in the medical record. Modifiers may be used to provide additional detail when billing for the EIP to the DME MAC. ⁸ Note: Drug administration codes may get billed to the MAC and the E-codes may get billed separately to the DME MAC. If the office bills the G-code to the MAC, the cost of the pump and supplies is bundled and should not be billed separately to the DME MAC. ¹³ Report any supplies as necessary.

^{*}Reporting policies for discarded units for payers other than traditional fee-for-service Medicare may vary; providers should check with their specific plans about policies related to billing for discarded drug and use of the JW and JZ modifiers.

Sample CMS-1500 Form: Physician Office Administration

APPROVED BY NAT PICA 1. MEDICARE (Medicareit)	(Medicaid#) (ID	N COMMITTEE (NUC	CHAMPVA (Member ID#) 3. PA 6. PA Se STATE 8. RE	_	H DATE M IONSHIP TO IN	SEX	1a. INSURED'S 4. INSURED'S A 7. INSURED'S A CITY ZIP CODE	JAME (Last Nar	me, First Name, Mi	PICA [[For Program in Item 1]] iddle Initial) STATE Include Area Code)	1.	
			a. EN		F?	PLACE (State)	a. INSURED'S C	DATE OF BIRTH DD YY M ID (Designate	М	SEX F		
NDC (BOX 24A SHADED AF payers (eg, Medicaid) m the HCPCS J-code. When qualifier "N4," indicating the 11-digit format. The u 3 spaces after the NDC, the payer-specific claim	ay require listir n required by the that an NDC fo nit of measure as UNX (X = the	orting BLINC ng the NDC i he payer, rep ollows, and t can also be e number of quirements	CYTO®, sor n addition ort the NI he NDC in reported, vials). Ver	to DC n =	appropr ICD-10-0 • Z51.1 immu • C91.0 leuke remis • C91.0 leuke	2 Encounter unotherapy 0 Acute ly emia not he ession OR 11 Acute ly emia, in rel 2 Acute ly emia, in rel des depen	osis code; er for antir y AND imphoblas aving achi imphoblas mission OF imphoblas lapse	eg, neoplasti tic eved tic tic cal recor	C ONS SI dersigne KIN CUI TO D TO CL TO D I L CL L CL	N7 GNATURE Lauthorize GNATURE Lauthorize d physician or supplier for GRENT OCCUPATION MM DD YY JURIERT SERVICES MM DD YY JAGNOSIS POINTER nter the letter (A—I) the diagnosis in I	L) that c	
1 N45551301600 MM DD YY 2 N45551301600 MM DD YY 3 MM DD YY	J.	PLACE OF	K. L. D. PROCEDURE: (Explain Unu CPT/HCPCS) J9039 J9039 96419	sual Circumstar	L. L OR SUPPLIES	DIAGN DSIS POIN FER A B A B	F. \$ CHARGES XXX XXX XXX	хх х	H. I. PSCI ID. Pen GUAL. NPI NPI NPI	UNITS (Box 24G) Report units of units administe discarded drug reported as 1 u 1 unit for initiat or refill of EIP	service ered and BLINC nit per r	d amount of YTO® dose ncg. Report
PLACE OF SERVICE (Box 24B Enter the appropriate 2-dig service code that correspondent of the control ocation where services are 11 Physician office	nit place of nds to the	N 20	Use the CF 96416 Che chemothe OR 96521 Refi OR G0498 Che setting us (eg, home includes fo	PT code remotherapy infusions infusi	epresenti py admini sion (mor maintena py admin e/clinic pu ary, rest l office/clii	ng the prostration, in the than 8 h ance of po istration, I imp/suppl nome, or a nic visit at	ocedure pentravenou ours), requ rtable pun V infusion ies, with c ssisted liv the conclu	erformed, s infusion uiring use np techniquontinuatiing) usinusion of t	such as in technique e of a porta ue; initiatio ion of the in g a portable he infusior	Iministration procitiation OR refill; e; initiation of prolible or implantable or implantable or implantable or implantable or infusion in the comple pump provided to be claims for drug	g, onged e pump e office/ imunity by the c	setting office/clinic;

This sample form is intended as a reference for coding and billing for product and associated services. It is not intended to be directive; the use of the recommended codes does not guarantee reimbursement. Healthcare providers may deem other codes or policies more appropriate and should select the coding options that most accurately reflect their internal system guidelines, payer requirements, practice patterns, and the services rendered. Healthcare providers are responsible for ensuring the accuracy and validity of all billing and claims for appropriate reimbursement.



Home Infusion - Multiple Payers (Medicare and Non-Medicare)

Item	Coding Information (ICD-10-CM ⁷ /CPT ⁹ /HCPCS ⁸ /NDC ¹¹)	Notes
Diagnosis: Encounter for drug therapy and ALL	Z51.12 Encounter for antineoplastic immunotherapy AND C91.00 Acute lymphoblastic leukemia not having achieved remission OR C91.01 Acute lymphoblastic leukemia, in remission OR C91.02 Acute lymphoblastic leukemia, in relapse	Report the appropriate ICD-10-CM code(s) to describe the patient's condition.
Procedure: Administration via CIVI using an EIP	G0090 Professional services, initial visit, for the administration of intravenous chemotherapy or other highly complex infusion drug or biological for each infusion drug administration calendar day in the individual's home, each 15 minutes! G0070 Professional services for the administration of intravenous chemotherapy or other intravenous highly complex drug or biological infusion for each infusion drug administration calendar day in the individual's home, each 15 minutes! 99601 Home infusion/specialty drug administration, per visit (up to 2 hours) 99602 Each additional hour S9329 Home infusion therapy, chemotherapy infusion; administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem S9330 Home infusion therapy, continuous (24 hours or more) chemotherapy infusion; administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem S9338 Home infusion therapy, immunotherapy, administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem S9379 Home infusion therapy, infusion therapy, not otherwise classified; administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem	Home infusion therapy services for Medicare beneficiaries receiving BLINCYTO® should be billed using G0090 for an initial visit and G0070 for subsequent visits. Some or all Medicare contractors may reject chemotherapy CPT codes with the availability of G0070 and G0090. These services must be reported to the A/B MAC, and are reimbursed by Medicare at rates set by the Medicare Physician Fee Schedule. They are billed and paid separately from the external infusion pump and drug, which are billed to the DME MAC and reimbursed under the DMEPOS Fee Schedule. Medicare requires that a claim for BLINCYTO® be billed no more than 30 days prior to the visit. Otherwise, payment for the home infusion therapy service will be denied.¹ These services may be covered by Medicaid, commercial plans, or Medicare Advantage plans.²0 CPT codes 99601 and 99602, as well as certain S-codes, may be used to report home infusion therapy services to other payer types other than FFS Medicare. Please note that FFS Medicare does not recognize S-codes, although other payers might.²0
Drug: BLINCYTO®	J9039 Injection, blinatumomab, 1 mcg JW Discarded drug/not administered to any patient JZ Zero drug amount discarded/not administered to any patient	Medicare requires that claims for BLINCYTO®, the pump, and supplies be sent to the DME MACs. Claims for home infusion therapy services must now be submitted separately and are processed by Part A/B MACs.¹ Medicare sets maximum utilization at 875 units of service (UOS), which is equivalent to 25 vials per month in this site of care.²¹ Many payers require the use of the modifier JW and JZ, which provides payment for the amount of drug or biologic discarded, as well as for the dose administered, up to the amount of the drug or biologic as indicated on the vial or label for an SDV.¹⁵ Note: Effective for dates of service on or after July 1, 2023, Medicare Part B claims require the use of the new JZ modifier for single-use vials and containers when there are no discarded drug amounts. Medicare claims also continue to require the use of the JW modifier (Drug amount discarded/not administered to any patient) for drugs and biologicals that are separately payable under Medicare Part B with discarded amounts from single-dose containers.*
	NDC: 55513016001 BLINCYTO® 35 mcg lyophilized powder, SDV IV solution stabilizer, 10 mL SDV	Some payers (eg, Medicaid) may require listing the NDC in addition to the HCPCS J-code. When reporting the NDC on claims, use the 11-digit NDC in the 5-4-2 format. Insert a leading zero in the appropriate section to complete the 5-4-2 digit format. Remove the dashes prior to entering the NDC on the claim form.
DME: EIP and supplies	E0779 Ambulatory infusion pump, mechanical, reusable, for infusion 8 hours or greater ²² E0781 Ambulatory infusion pump, single or multiple channels, electric or battery operated, with administrative equipment, worn by patient ²² A4222 Infusion supplies for external drug infusion pump, per cassette or bag Modifiers for EIP -KD Drug or biologic infused through DME -RR Rental -KH DMEPOS item, initial claim or first rental month -KI DMEPOS item, second or third rental months -KJ DMEPOS item, fourth to 15th rental months	Report the appropriate EIP code and appropriate modifier(s) as documented in the medical record. Modifiers may be used to provide additional detail when billing for the EIP to the DME MAC.8 Report any supplies as necessary.

^{*}Reporting policies for discarded units for payers other than traditional fee-for-service Medicare may vary; providers should check with their specific plans about policies related to billing for discarded drug and use of the JW and JZ modifiers.

BLINCYTO®

Billing Information Sheet



Sample CMS-1500 Form: Medicare DME MAC for BLINCYTO®, Pump, and Related Supplies by DME Supplier

HEALTH INSURANCE CLAIM FOI APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE IN		CARRIER +
MEDICARE MEDICAID TRICARE (Medicare#) (Medicaid#) (ID#/DoD#) 2. PATIENT'S NAME (Last Name, First Name, Middle Initial)	CHAMPVA GROUP FECA BIX (LING OTHER (Member IDH)	1a. INSURED'S I.D. NUMBER (For Program in Item 1) 4. INSURED'S NAME (Last Name, First Name, Middle Initial)
sample claim form on page 11 for guidance on	6. PATIENT RELATIONSHIP TO INSURED Self Spouse Child Other STATE 8. RESERVED FOR NUCC USE Code) 10. IS PATIENT'S CONDITION RELATED TO:	7. INSURED'S ADDRESS (No., Street) CITY ZIP CODE TELEPHONE (Include Area Code) () 11. INSURED'S POLICY GROUP OR FECA NUMBER
b. RESERVED FOR NUCC USE	a. EMPLOYMENT? (Current or Previous) YES NO b. AUTO ACCIDENT? PLACE (State)	a. INSURED'S DATE OF BIRTH MM DD YY M F SEX b. OTHER CLAIM ID (Designated by NUCC)
NDC (BOX 24A SHADED AREA): When required by the payer, report the NDC qualifier "N4," indicating that an NDC follows, and the NDC in the 11-digit format. The unit of measure can also be reported, 3 spaces after the NDC, as UNX (X = the number of vials)	c. OTHER ACCIDENT? YES NO 10d. CLAIM CODES (Designated by NUCC) DMPLETING & SIGNING THIS FORM. Ultraze the release of any medical or other information necessary neffic either to myself or to the party who accepts assignment DATE	c. INSURANCE PLAN NAME OR PROGRAM NAME d. IS THERE ANOTHER HEALTH BENEFIT PLAN? YES NO If yes, complete items 9, 9a, and 9d, 13. INSURED'S OR AUTHORIZED PERSON'S SIGNATURE Lauthorize payment of medical benefits to the undersigned physician or supplier for services described below. SIGNED
1. DATE OF CURRENT ILLNESS. INJURY, or PREGNANCY OUAL. 1. NAME OF REFERRING PROVIDER OR OTHER SOURCE 1. ADDITIONAL CLAIM INFORMATION (Designated by NUCC 2. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relat. 2. Z51.12 B. C91.02 F. L	DIAGNOSIS (BOX 21): Enter the appropriate diagnosis Z51.12 Encounter for antineoplas C91.04 Acute lymphoblastic leuk	stic immunotherapy AND kemia not having achieved remission OR kemia, in remission OR kemia, in relapse
2 A. DATE(S) OF SERVICE B. C. PLACE OF SERVICE EMG 1 N455513016001 UNX MM DD YY MM DD YY 12	D. PROCEDURES, SERVICES, OR SUPPLIES (Explain Unusual Circumstances) CPT/HCPCS MODIFIER J9039 A B	F. DAYS LANGES SCHARGES SCHARGES WITS REPORT UNITS (Box 24G): Report units of service for both units administered and amount units administered and amou
2 N455513016001 UNX MM DD YY MM DD YY 12 3 3 MM DD YY MM DD YY 12	J9039 JW AB E0781 RR KH AB	of discarded drug. BLINCYTO® dose reported as 1 unit per mcg. XXX XX X 1 NPI XXX XX 1 NPI NPI TOTAL TOTA
4 MM DD YY MM DD YY 12 5	A4222 A B	DIAGNOSIS POINTER (Box 24E): Enter the letter (A–L) that corresponds to the diagnosis in Box 21
PLACE OF SERVICE (Box 24B): Enter the appropriate 2-digit place of service code that corresponds to the location where services are rendered; eg, • 12 Home SIGNED DATE	JW/JZ Discard Modifier: JW or JZ mo DME external infusion pump claims single-use containers IV Pump: E0781 Ambulatory infus A4222 Infusion supplies for extern cassette or infusion option	odes and modifiers; eg, difier required for Medicare including infused drugs in sion pump

This sample form is intended as a reference for coding and billing for product and associated services. It is not intended to be directive; the use of the recommended codes does not guarantee reimbursement. Healthcare providers may deem other codes or policies more appropriate and should select the coding options that most accurately reflect their internal system guidelines, payer requirements, practice patterns, and the services rendered. Healthcare providers are responsible for ensuring the accuracy and validity of all billing and claims for appropriate reimbursement.

Sample CMS-1500 form: Medicare A/B MAC for Home Infusion Therapy Services by Home Infusion Therapy Supplier

	1. MEDICARE MEDICAID TRICARE CHAMPV (Medicare#) (Medicaid#) (ID#/DoD#) (Member I 2. PATIENT'S NAME (Last Name, First Name, Middle Initial)	MENTUDIAN DIVING	A Insured's I.D. Number (For Program In Item 1) 4. Insured's Name (Last Name, First Name, Middle Initial)
ome infus rnished a etting. Thi r billing h edicare p age 10 for	dicare requires separate claims for ion therapy services and for drugs s items of DME in the home infusion s sample claim shows an example ome infusion therapy services for a atient. See the sample claim form on guidance on billing for drugs furnished of DME for a Medicare beneficiary	6. PATIENT RELATIONSHIP TO INSURED Self Spouse Child Other 8. RESERVED FOR NUCC USE	7. INSURED'S ADDRESS (No., Street) CITY ZIP CODE TELEPHONE (Include Area Code) () 11. INSURED'S POLICY GROUP OR FECA NUMBER a. INSURED'S DATE OF BIRTH MM DD YY M F b. OTHER CLAIM ID (Designated by NUCC)
	a, OTHER INSURED'S POLICY OR GROUP NUMBER	a. EMPLOYMENT? (Current or Previous) YES NO	a. INSURED'S DATE OF BIRTH SEX
	b. RESERVED FOR NUCC USE	b. AUTO ACCIDENT? PLACE (State)	b. OTHER CLAIM ID (Designated by NUCC)
	c. RESERVED FOR NUCC USE	c. OTHER ACCIDENT?	d. IS THERE ANOTHER HEALTH BENEFIT PLAN?
	d. INSURANCE PLAN NAME OR PROGRAM NAME	10d. CLAIM CODES (Designated by NUCC)	
	READ BACK OF FORM BEFORE COMPLETIN 12. PATIENT'S OR AUTHORIZED PERSON'S SIGNATURE I authorize the to process this claim. I also request payment of government benefits either below.	release of any medical or other information necessary	YES NO If yes, complete items 9, 9a, and 9d.
	SIGNED	DATEOTHER DATE	SIGNED
	17. NAME OF REFERRING PROVIDER OR OTHER SOURCE 19. ADDITIONAL CLAIM INFORMATION (Designated by NUCC) 21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate A-L to s A. Z51.12 B. C91.02	Enter the appropriate diagnosis Z51.12 Encounter for antineoplas C91.00 Acute lymphoblastic leuk C91.01 Acute lymphoblastic leuk C91.02 Acute lymphoblastic leuk Final codes depend on medical	stic immunotherapy AND kemia not having achieved remission OR kemia, in remission OR kemia, in relapse
	E. F. G. K. [L. L.	
	24. A. DATE(S) OF SERVICE B. C. D. PROCE From To PLACE OF (Explication of the control of the contr	EDURES, SERVICES, OR SUPPLIES ain Unusual Circumstances) PCS MODIFIER POINTER	S CHARGES UNITS (Box 24G): Report units of service
	MM DD YY MM DD YY 12 G0090	D A B	XXX XX X NPI for the administration BLINCYTO®, reported a
	2 MM DD YY MM DD YY 12 J9039	A B	0 00 X NPI 1 unit per 15 minutes o
	3		NPI S
	ERVICE (Box 24B): propriate 2-digit place of service porresponds to the location where	DIAGNOSIS POINTER letter (A–L) that coldiagnosis in Box 21	NPI NPI
le that c	e rendered; eg,		NPI G

This sample form is intended as a reference for coding and billing for product and associated services. It is not intended to be directive; the use of the recommended codes does not guarantee reimbursement. Healthcare providers may deem other codes or policies more appropriate and should select the coding options that most accurately reflect their internal system guidelines, payer requirements, practice patterns, and the services rendered. Healthcare providers are responsible for ensuring the accuracy and validity of all billing and claims for appropriate reimbursement.

BLINCYTO® (blinatumomab) for (blinatumomab) for single-use vial

13

Sample CMS-1500 Form: Non-Medicare Payer by Home Infusion Provider

	APPROVED BY NATIONAL UNIFORM CLA	, , , , ,				R 1a. INSURED'S I.D	NUMBER		PICA PICA
		TRICARE CHAMP (ID#/DoD#) (Member	HE/	OUP ALTH PLAN BI (II	ECA OTHER BLK LUNG (ID#)	1a. INSURED'S I.L	. NUMBER	(For Progran	m in Item 1)
	2. PATIENT'S NAME (Last Name, First Nat	me, Middle Initial)	3. PATIENT	T'S BIRTH DATE	SEX F	4. INSURED'S NAI	ME (Last Nan	ne, First Name, Middle Initial)	
	5. PATIENT'S ADDRESS (No., Street)		_	FRELATIONSHIP T		7. INSURED'S ADI	DRESS (No.,	Street)	
	СПҮ	STATE	Self 8. RESERV	Spouse Child ED FOR NUCC US		CITY			STATE
	ZIP CODE TELEPH	HONE (Include Area Code)				ZIP CODE		TELEPHONE (Include Area	a Code)
	()				November 1	LIOV OBSULT	()	
	9. OTHER INSURED'S NAME (Last Name,	, First Name, Middle Initial)	10. IS PATI	ENT'S CONDITION	N RELATED TO:			P OR FECA NUMBER	
	a. OTHER INSURED'S POLICY OR GROU	JP NUMBER	a. EMPLOY	MENT? (Current or	or Previous)	a. INSURED'S DA	E OF BIRTH	SEX M	F
	b. RESERVED FOR NUCC USE		b. AUTO A	CCIDENT?	PLACE (State)	b. OTHER CLAIM	D (Designate	ed by NUCC)	
	c. RESERVED FOR NUCC USE		c. OTHER	YES ACCIDENT?	NO	c. INSURANCE PL	AN NAME OF	R PROGRAM NAME	
	d, INSURANCE PLAN NAME OR PROGRA	AM NAME	10d CLAIN	YES CODES (Designat	NO ted by NUICC)	d IS THERE AND	THER HEALT	'H BENEFIT PLAN?	
					nod by Noce /	YES	NO	If yes, complete items 9, 9a,	
	12. PATIENT'S OR AUTHORIZED PERSO to process this claim. Lalso request payor	F FORM BEFORE COMPLETINDN'S SIGNATURE I authorize the ment of government benefits either	e release of any	i THIS FORM. y medical or other in o the party who acce	nformation necessary epts assignment		dical benefits	ED PERSON'S SIGNATURE I to the undersigned physician	
follows, a measure	24A SHADED AREA): When reque NDC qualifier "N4," indicating and the NDC in the 11-digit form: can also be reported, 3 spaces the number of vials). Verify the	g that an NDC eat. The unit of after the NDC, as	_ D	MM D	IS (BOX 21): Er	FROM		TO WORK IN CURRENT OCC	
follows, a measure UNX (X =	e NDC qualifier "N4," indicating and the NDC in the 11-digit forms can also be reported, 3 spaces the number of vials). Verify the omission requirements	g that an NDC lat. The unit of a after the NDC, as e payer-specific	HER DAT	DIAGNOSI	IS (BOX 21): Er Encounter fo Acute lymph	nter the appror antineoplastic leulhoblastic leulhoblastic leulhoblastic leunhoblastic leunhoblastic leulhoblastic	opriate o stic imm kemia no kemia, i kemia, i	diagnosis code; eq nunotherapy AND ot having achieved n remission OR n relapse	g, ICD-10-CM:
follows, a measure UNX (X =	e NDC qualifier "N4," indicating and the NDC in the 11-digit forms can also be reported, 3 spaces the number of vials). Verify the omission requirements 21. DIAGNOSIS OR NATURE OF ILLNESS A. LZ51.12 B. C. E. L. F. L.	g that an NDC nat. The unit of s after the NDC, as e payer-specific S OR INJURY Relate A-L to ser 191.02 C.1 G.1	NPI rvice line belov	DIAGNOSI	IS (BOX 21): Er E Encounter fo D Acute lymph 1 Acute lymph 2 Acute lymph des depend o	nter the appror antineoplastic leulhoblastic leulhoblastic leulhoblastic leunhoblastic leunhoblastic leulhoblastic	opriate o stic imm kemia no kemia, i kemia, i	diagnosis code; eq nunotherapy AND ot having achieved n remission OR n relapse	g, ICD-10-CM:
follows, a measure UNX (X =	e NDC qualifier "N4," indicating and the NDC in the 11-digit forms can also be reported, 3 spaces the number of vials). Verify the omission requirements 21. DIAGNOSIS OR NATURE OF ILLNESS A. Z51.12 E. L I. L 24. A. DATE(S) OF SERVICE From To	g that an NDC nat. The unit of s after the NDC, as e payer-specific S OR INJURY Relate A-L to ser C91.02 G. I B. C. D. PROCC PLACE OF CEXPLACE O	NPI VICE line below	DIAGNOSI	IS (BOX 21): Er P. Encounter for D. Acute lymph 1 Acute lymph 2 Acute lymph des depend o	nter the appror antineoplastic leul hoblastic leul n medical re	opriate ostic imm kemia no kemia, i kemia, i cord doo	diagnosis code; eq nunotherapy AND ot having achieved n remission OR n relapse cumentation	g, ICD-10-CM: d remission 0R
follows, a measure UNX (X =	e NDC qualifier "N4," indicating and the NDC in the 11-digit form can also be reported, 3 spaces the number of vials). Verify the omission requirements 21. DIAGNOSIS OR NATURE OF ILLNESS A. Z51.12 E. L. J. L. J. L. 24. A. DATE(S) OF SERVICE MM DD YY MM DD YY N455513016001 UNX	g that an NDC lat. The unit of later the NDC, as	NPI VICE line below	DIAGNOSI	IS (BOX 21): Er Encounter fo D Acute lymph 1 Acute lymph 2 Acute lymph des depend o	nter the appror antineoplastic leuloblastic	opriate (stic imm kemia n kemia, i kemia, ii cord doo	diagnosis code; equunotherapy AND ot having achieved necession OR necession of the properties of the p	g, ICD-10-CM: d remission OR TER (Box 24E): Enter the torresponds to the
follows, a measure UNX (X =	e NDC qualifier "N4," indicating and the NDC in the 11-digit form can also be reported, 3 spaces the number of vials). Verify the omission requirements 21. DIAGNOSIS OR NATURE OF ILLNESS A. Z51.12 B. C E. L F. F. L 1 L DATE(S) OF SERVICE From DD YY MM DD YY	g that an NDC lat. The unit of later the NDC, as	HER DAT	DIAGNOSI	IS (BOX 21): Er P. Encounter for D. Acute lymph 1 Acute lymph 2 Acute lymph des depend o	nter the appror antineoplastic leul hoblastic leul n medical re	opriate (stic imm kemia n kemia, i kemia, ii cord doo	diagnosis code; equunotherapy AND ot having achieved necession OR necession or relapse cumentation	g, ICD-10-CM: d remission OR TER (Box 24E): Enter the torresponds to the
follows, a measure UNX (X =	e NDC qualifier "N4," indicating and the NDC in the 11-digit forms can also be reported, 3 spaces the number of vials). Verify the omission requirements 21. DIAGNOSIS OR NATURE OF ILLNESS A. Z51.12 E	g that an NDC lat. The unit of safter the NDC, as e payer-specific S OR INJURY Relate A-L to see 191.02 C. I B. C. PLACE OF LY SERVICE EMG CPT/HCI Y 12 J9039	NPI NPI EDURES, SEI lain Unusual OPCS 9	DIAGNOSI Z51.12 C91.00 G91.01 C91.02 Final coc H. RWICES, OR SUPP ircumstances) MODIFIER	IS (BOX 21): Er P. Encounter for D. Acute lymph 1. Acute lymph 2. Acute lymph des depend o	nter the appror antineoplastic leuloblastic	opriate ostic imm kemia nokemia, i kemia, i kemia, in cord doc	diagnosis code; equunotherapy AND ot having achieved necession OR necession of the properties of the p	TER (Box 24E): Enter the torresponds to the
follows, a measure UNX (X =	e NDC qualifier "N4," indicating and the NDC in the 11-digit forms can also be reported, 3 spaces the number of vials). Verify the omission requirements 21. DIAGNOSIS OR NATURE OF ILLNESS A. Z51.12 B. C E. L. J. J. L. 24. A. DATE(S) OF SERVICE MM DD YY MM DD YY N455513016001 UNX MM DD YY MM DD YY 2 N455513016001 UNX	that an NDC hat. The unit of safter the NDC, as a payer-specific sort in the safter the NDC, as a payer-specific sort in the safter the NDC, as a payer-specific sort in the safter the NDC, as a payer-specific sort in the safter the NDC, as a payer-specific sort in the safter the NDC, as a payer specific sort in the NDC, as a payer specific sort in the NDC, as a payer specific sort in the NDC, as a payer specifi	NPI NPI NPI Privice line below L L L EBURES, SEI Idain Unusual C PCS 9 JV	DIAGNOSI Z51.12 C91.00 G91.01 C91.02 Final coc H. RWICES, OR SUPP ircumstances) MODIFIER	IS (BOX 21): Er P. Encounter for D. Acute lymph 1 Acute lymph 2 Acute lymph des depend o	nter the appror antineoplastic leulhoblastic	opriate of stic immediate of s	diagnosis code; equunotherapy AND obt having achieved in remission OR in reliapse cumentation DIAGNOSIS POIN letter (A-L) that diagnosis in Boy UNITS (Box 24 Report units)	TER (Box 24E): Enter the torresponds to the x 21 1G): of service for both units
follows, a measure UNX (X =	e NDC qualifier "N4," indicating and the NDC in the 11-digit form: can also be reported, 3 spaces the number of vials). Verify the omission requirements 21. DIAGNOSIS OR NATURE OF ILLNESS A. Z51.12 E. L. F. L. 1. L. J. J. L. 24. A. DATE(S) OF SERVICE From DD YY MM DD YY NA55513016001 UNX MM DD YY MM DD YY N455513016001 UNX MM DD YY MM DD YY 3 MM DD YY MM DD YY 3 MM DD YY MM DD YY	that an NDC hat. The unit of safter the NDC, as a payer-specific so the safter the NDC, as a payer-specific so	NPI	DIAGNOSI Z51.12 C91.00 G91.01 C91.02 Final coc H. RWICES, OR SUPP ircumstances) MODIFIER	IS (BOX 21): Er P. E. E. C.	nter the appror antineoplastic leul hoblastic leul noblastic leul nedical research	opriate ostic immersione opriate ostic immersione opriate ostic immersione oprimalista opr	diagnosis code; economotion of the vine of the vertical of the	g, ICD-10-CM: d remission OR TER (Box 24E): Enter the t corresponds to the x 21 IG): of service for both units nistered and amount drug. BLINCYTO® dose
follows, a measure UNX (X =	e NDC qualifier "N4," indicating and the NDC in the 11-digit form: can also be reported, 3 spaces the number of vials). Verify the pmission requirements 21. DIAGNOSIS OR NATURE OF ILLNESS A. Z51.12 E. L. J. L. J. L. 24. A. DATE(S) OF SERVICE From To MM DD YY MM DD YY 1 N455513016001 UNX MM DD YY MM DD YY 2 N455533016001 UNX MM DD YY MM DD YY 3 MM DD YY MM DD YY 3 MM DD YY MM DD YY 3 MM DD YY MM DD YY	that an NDC hat. The unit of safter the NDC, as a payer-specific so the safter the NDC, as a payer-specific so	NPI	DIAGNOSI Z51.12 C91.00 G91.01 C91.02 Final coc H. RWICES, OR SUPP ircumstances) MODIFIER	IS (BOX 21): Er Encounter for D Acute lymph 1 Acute lymph des depend o DIAGNOSIS POI TER A B	nter the appror antineoplastic leul hoblastic leul noblastic leul nedical research	opriate of stic immediate of s	diagnosis code; economotion of the vine of the vertical of the	g, ICD-10-CM: d remission 0R TER (Box 24E): Enter the t corresponds to the x 21 1G): of service for both units nistered and amount

This sample form is intended as a reference for coding and billing for product and associated services. It is not intended to be directive; the use of the recommended codes does not guarantee reimbursement. Healthcare providers may deem other codes or policies more appropriate and should select the coding options that most accurately reflect their internal system guidelines, payer requirements, practice patterns, and the services rendered. Healthcare providers are responsible for ensuring the accuracy and validity of all billing and claims for appropriate reimbursement.

BLINCYTO® Dosing Options¹¹

Dosing option	Dose per vial X number of SDVs*	Number of billing units
24-hour	35 mcg X 1 vial	35
48-hour	35 mcg X 1-2 vials	35-70
7-day	35 mcg X 4-6 vials	140-210

^{*}Number of SDVs depends on dose, infusion duration, and patient's weight.¹¹

Key Considerations for the BLINCYTO® 7-day Infusion Option (7-DIO)



Minor variations are expected in coding, billing, and claims filing for the BLINCYTO® 7-DIO.²⁰



The 7-DIO requires 6 vials of BLINCYTO® and 1 vial of IV Solution Stabilizer for patients \geq 45 kg. For patients weighing less than 45 kg, 4 to 5 vials are required. The safety of the administration of BLINCYTO® at a BSA of less than 0.4 m² has not been established.¹¹ Refer to the Prescribing Information for details on handling and preparation.



If the units field on a claim form cannot accommodate more than 99 units, utilize multiple lines to capture all units (eg, 99+98+13). Payers may require separate reporting of drug units administered and discarded.²⁰



Less frequent claim submissions are expected with utilization of the 7-DIO. Typically the entire 7-DIO can be billed on the day of administration/refill. However, be sure to refer to payer guidelines for maximum daily quantity limits. Apply the appropriate dates of service as needed.²⁰



If the 7-DIO is interrupted mid-treatment, refer to payer guidelines for reporting and documentation in these cases. If full reimbursement is withheld by the payer, refer to Amgen's Product Return Policy for assistance.



Existing codes and modifiers are adequate to report BLINCYTO® and its related services; however, payer requirements may vary with respect to:²⁰

- The entities that can bill for DME and the associated supplies
- The number of units billed for BLINCYTO® J9039 (HCPCS units vs number of vials)
- Covered diagnosis codes
- Covered nursing services (eg, infusion services at patient's home)
- Drug claim submission options (eg, 1 or more dates of service on claims)
- Reporting policies for discarded units for payers other than traditional fee-for-service Medicare
 may vary; providers should check with their specific plans about policies related to billing for
 discarded drug and use of the JW and JZ modifiers.

UNDERSTANDING EXAMPLES OF



REIMBURSEMENT ACROSS SITES OF CARE

A BLINCYTO® patient transitions through multiple sites of care. This guide shows how major payers in the United States (commercial plans, Medicare, and Medicaid) offer coverage in each setting and reimburse for each component of care:



Drug







Pump and Supplies Hospitalization

Professional Services (ie, drug administration)

INDICATIONS

BLINCYTO® (blinatumomab) is indicated for the treatment of CD19-positive B-cell precursor acute lymphoblastic leukemia (ALL) in adult and pediatric patients one month and older with:

- Philadelphia chromosome-negative disease in the consolidation phase of multiphase chemotherapy
- Minimal residual disease (MRD) greater than or equal to 0.1% in first or second complete remission
- Relapsed or refractory disease

IMPORTANT SAFETY INFORMATION


WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGICAL TOXICITIES including IMMUNE EFFECTOR **CELL-ASSOCIATED NEUROTOXICITY SYNDROME**

- · Cytokine Release Syndrome (CRS), which may be life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® and treat with corticosteroids as recommended.
- Neurological toxicities, including immune effector cell-associated neurotoxicity syndrome (ICANS) which may be severe, life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® as recommended.

Please see additional Important Safety Information for BLINCYTO® on pages 18-19.

BLINCYTO® (blinatumomab) Reimbursement Process

Coverage of BLINCYTO® and its administration is required in all these sites of care to avoid interruption in treatment.



The scenarios depicted above illustrate the most common ones for accessing BLINCYTO® via the buy-and-bill acquisition process, where the entity that acquires the product also administers it to the patient.

BLINCYTO® can also be acquired via a specialty pharmacy provider, including:

- Third-party specialty pharmacies that contract with a payer to supply specialty products covered under the medical benefit
- Specialty pharmacies owned by hospitals, physician offices, ambulatory infusion clinics, and/or home infusion companies that may also administer the medication

BLINCYTO® Reimbursement Across Transitions in Site of Care

BLINCYTO®-eligible patients need coverage for the following: Drug + Pump + Hospitalization + Administration

Inpatient Hospit	al			
Components of BLINCYTO® Care	Commercially Insured Patients	FFS Medicare Patients	FFS Medicaid Patients	
BLINCYTO®	MS-DRG-based or global payment; typically includes BLINCYTO®23	MS-DRG payment includes BLINCYTO® ²⁴	APR-DRG-based payment; typically includes BLINCYTO®27	
	Reimbursement varies by contracts between providers and payers	Covered under Medicare Part A benefit ²⁵	Reimbursement varies by state; may follow Medicare	
Pump and Supplies	Some hospitals, in their contracts with managed care organizations, may negotiate a "carve out" benefit for drugs such as BLINCYTO® • May allow separate payment of	Hospital may be eligible for outlier payments if cost of admission exceeds certain threshold	allowable amounts	
Hospitalization	such drugs outside of the bundled payment for inpatient services	Reimbursement varies for the 11 IPPS-Exempt Cancer Hospitals ²⁶		
Professional Services	Physician services may be covered separately outside of the bundled payment	Physician services may be covered and reimbursed according to the MPFS under Medicare Part B benefit	Physician services may be covered and paid outside of the bundled payment	

Key: APR-DRG-All Patient-Refined Diagnosis Related Groups; FFS-fee-for-service; IPPS-Inpatient Prospective Payment System; MPFS – Medicare Physician Fee Schedule; MS-DRG – Medicare Severity Diagnosis-Related Group.

Outpatient Hosp	ital		
Components of BLINCYTO® Care	Commercially Insured Patients	FFS Medicare Patients	FFS Medicaid Patients
BLINCYTO®	Reimbursed based on contracted rates; methodology varies Examples: ²⁸ • ASP + X% • WAC + X% • AWP-X% May need prior authorization	Covered under Medicare Part B benefit Typically reimbursed based on ASP + 6% when administered in a 340B hospital setting (with 2% sequestration reduction) ^{29,30} MUE cap of 210 units (approx. 6 vials) per date of service applies ^{31,8}	Reimbursement may be similar to Medicare OR State-defined limit (eg, California uses a federal upper limit) 32 May need prior authorization
Pump and Supplies	Reimbursement is bundled into the payment for the infusion service	Covered under Medicare Part B benefit Reimbursement is bundled into the payment for the infusion service	Reimbursed based on fee schedule or bundled into the payment for the infusion service Rates vary by state
Professional Services	Reimbursed based on contracted rate	Reimbursed based on the Medicare OPPS	

Key: ASP-average sales price; AWP-average wholesale price; FFS-fee-for-service; MUE-medically unlikely edit; OPPS-Outpatient Prospective Payment System;

Note: The information here describes coverage and payment for BLINCYTO® under FFS Medicare and FFS Medicaid. Coverage and payment for patients enrolled in Medicare Advantage and/or Medicaid managed care organizations varies widely and is often similar to commercial insurance.

Physician Office					
Components of BLINCYTO® Care	Commercially Insured Patients	FFS Medicare Patients	FFS Medicaid Patients		
BLINCYTO®	Reimbursed based on contracted rates; methodology varies Examples: ²⁸ • ASP + X% • WAC + X% • AWP-X% May need prior authorization	Covered under Medicare Part B benefit Typically reimbursed based on ASP + 6% (with 2% sequestration reduction) ^{30,33} MUE cap of 210 units (approx. 6 vials) per date of service applies ^{8,19}	Reimbursement may be similar to Medicare OR State-defined limit (eg, California uses a federal upper limit) ³² May need prior authorization		
Pump and Supplies	Reimbursed based on contracted rate and bundled into payment for the infusion service	Covered under Medicare Part B benefit Reimbursement is bundled into the payment for the infusion service	Typically reimbursed based on fee schedule or bundled into the payment for the infusion service Rates vary by state		
Professional Services	Reimbursed based on contracted rate	Reimbursed based on the MPFS			

Key: ASP-average sales price; AWP-average wholesale price; FFS-fee-for-service; MPFS-Medicare Physician Fee Schedule; MUE-medically unlikely edit; WAC-wholesale acquisition cost.

Home Infusion			
Components of BLINCYTO® Care	Commercially Insured Patients	FFS Medicare Patients	FFS Medicaid Patients
BLINCYTO®	Reimbursed based on contracted rates; methodology varies Examples: ²⁸ • ASP + X% • WAC + X% • AWP – X% May need prior authorization	Covered under Medicare Part B as long as it is supplied in a covered external infusion pump and the IV is initiated in home infusion setting ³⁴ Typically reimbursed based on ASP + 6% (with 2% sequestration reduction) ^{30,33} Billing cap of 25 vials per month applies ²¹	Reimbursement may be similar to Medicare OR State-defined limit (eg, California uses a federal upper limit) ³⁷ May need prior authorization
Pump and Supplies	Reimbursed based on contracted rate	Covered under Medicare Part B benefit Reimbursed as part of the Medicare DMEPOS Fee Schedule ³⁵	Typically reimbursed based on fee schedule Rates vary by state
Professional Services	Reimbursed based on contracted rate	Covered under Part B Reimbursed under the home infusion therapy services benefit in 15-minute increments for applicable providers ³⁶	

 $Key: ASP-average \ sales \ price; AWP-average \ wholesale \ price; DMEPOS-Durable \ Medical \ Equipment \ Prosthetics, Orthotics, and \ Supplies; FFS-fee-for-service; \ DMEPOS-Durable \ Medical \ Equipment \ Prosthetics, Orthotics, and \ Supplies; FFS-fee-for-service; \ PFS-fee-for-service; \ PF$ WAC-wholesale acquisition cost.

Note: Medicare home infusion benefit is distinct and separate from the Medicare home health benefit.

BLINCYTO®

Indications and Important Safety Information



INDICATIONS

BLINCYTO® (blinatumomab) is indicated for the treatment of CD19-positive B-cell precursor acute lymphoblastic leukemia (ALL) in adult and pediatric patients one month and older with:

- Philadelphia chromosome-negative disease in the consolidation phase of multiphase chemotherapy
- · Minimal residual disease (MRD) greater than or equal to 0.1% in first or second complete remission
- Relapsed or refractory disease

IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGICAL TOXICITIES including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

- Cytokine Release Syndrome (CRS), which may be life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® and treat with corticosteroids as recommended.
- Neurological toxicities, including immune effector cell-associated neurotoxicity syndrome (ICANS) which may be severe, life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® as recommended.

Contraindications

BLINCYTO® is contraindicated in patients with a known hypersensitivity to blinatumomab or to any component of the product formulation.

Warnings and Precautions

- Cytokine Release Syndrome (CRS): CRS, which may be life-threatening or fatal, occurred in patients receiving BLINCYTO®. The median time to onset of CRS is 2 days after the start of infusion and the median time to resolution of CRS was 5 days among cases that resolved. Closely monitor and advise patients to contact their healthcare professional for signs and symptoms of serious adverse events such as fever, headache, nausea, asthenia, hypotension, increased alanine aminotransferase (ALT), increased aspartate aminotransferase (AST), increased total bilirubin, and disseminated intravascular coagulation (DIC). The manifestations of CRS after treatment with BLINCYTO® overlap with those of infusion reactions, capillary leak syndrome (CLS), and hemophagocytic histiocytosis/macrophage activation syndrome (MAS). Using all of these terms to define CRS in clinical trials of BLINCYTO, CRS was reported in 15% of patients with R/R ALL, in 7% of patients with MRD-positive ALL, and in 16% of patients receiving BLINCYTO® cycles in the consolidation phase of therapy. If severe CRS occurs, interrupt BLINCYTO® until CRS resolves. Discontinue BLINCYTO® permanently if life-threatening CRS occurs. Administer corticosteroids for severe or life-threatening CRS.
- Neurological Toxicities, including Immune Effector Cell-Associated Neurotoxicity Syndrome: BLINCYTO® can cause serious or life-threatening neurologic toxicity, including ICANS. The incidence of neurologic toxicities in clinical trials was approximately 65%. The median time to the first event was within the first 2 weeks of BLINCYTO® treatment. The most common (≥ 10%) manifestations of neurological toxicity were headache and tremor. Grade 3 or higher neurological toxicities occurred in approximately 13% of patients, including encephalopathy, convulsions, speech disorders, disturbances in consciousness, confusion and disorientation, and coordination and balance disorders. Manifestations of neurological toxicity included cranial nerve disorders. The majority of neurologic toxicities resolved following interruption of BLINCYTO®, but some resulted in treatment discontinuation.
 - The incidence of signs and symptoms consistent with ICANS in clinical trials was 7.5%. The onset of ICANS can be concurrent with CRS, following resolution of CRS, or in the absence of CRS. There is limited experience with BLINCYTO® in patients with active ALL in the central nervous system (CNS) or a history of neurologic events. Patients with a history or presence of clinically relevant CNS pathology were excluded from clinical studies. Patients with Down Syndrome over the age of 10 years may have a higher risk of seizures with BLINCYTO® therapy.
 - Monitor patients for signs and symptoms of neurological toxicities, including ICANS, and interrupt or discontinue BLINCYTO® as outlined in the PI. Advise outpatients to contact their healthcare professional if they develop signs or symptoms of neurological toxicities.
- Infections: Approximately 25% of patients receiving BLINCYTO® in clinical trials experienced serious infections such as sepsis, pneumonia, bacteremia, opportunistic infections, and catheter-site infections, some of which were life-threatening or fatal. Administer prophylactic antibiotics and employ surveillance testing as appropriate during treatment. Monitor patients for signs or symptoms of infection and treat appropriately, including interruption or discontinuation of BLINCYTO® as needed.
- Tumor Lysis Syndrome (TLS), which may be life-threatening or fatal, has been observed. Preventive measures, including pretreatment nontoxic cytoreduction and on-treatment hydration, should be used during BLINCYTO® treatment. Monitor patients for signs and symptoms of TLS and interrupt or discontinue BLINCYTO® as needed to manage these events.
- **Neutropenia and Febrile Neutropenia,** including life-threatening cases, have been observed. Monitor appropriate laboratory parameters (including, but not limited to, white blood cell count and absolute neutrophil count) during BLINCYTO® infusion and interrupt BLINCYTO® if prolonged neutropenia occurs.
- Effects on Ability to Drive and Use Machines: Due to the possibility of neurological events, including seizures and ICANS, patients receiving BLINCYTO® are at risk for loss of consciousness, and should be advised against driving and engaging in hazardous occupations or activities such as operating heavy or potentially dangerous machinery while BLINCYTO® is being administered.
- Elevated Liver Enzymes: Transient elevations in liver enzymes have been associated with BLINCYTO® treatment with a median time to onset of 3 days. In patients receiving BLINCYTO®, although the majority of these events were observed in the setting of CRS, some cases of elevated liver enzymes were observed outside the setting of CRS, with a median time to onset of 19 days. Grade 3 or greater elevations in liver enzymes occurred in approximately 7% of patients outside the setting of CRS and resulted in treatment discontinuation in less than

IMPORTANT SAFETY INFORMATION (continued)

1% of patients. Monitor ALT, AST, gamma-glutamyl transferase, and total blood bilirubin prior to the start of and during BLINCYTO® treatment. BLINCYTO® treatment should be interrupted if transaminases rise to > 5 times the upper limit of normal (ULN) or if total bilirubin rises to > 3 times ULN.

- Pancreatitis: Fatal pancreatitis has been reported in patients receiving BLINCYTO® in combination with dexamethasone in clinical trials and the post-marketing setting. Evaluate patients who develop signs and symptoms of pancreatitis and interrupt or discontinue BLINCYTO® and dexamethasone as needed.
- **Leukoencephalopathy:** Although the clinical significance is unknown, cranial magnetic resonance imaging (MRI) changes showing leukoencephalopathy have been observed in patients receiving BLINCYTO®, especially in patients previously treated with cranial irradiation and antileukemic chemotherapy.
- **Preparation and administration** errors have occurred with BLINCYTO® treatment. Follow instructions for preparation (including admixing) and administration in the PI strictly to minimize medication errors (including underdose and overdose).
- Immunization: Vaccination with live virus vaccines is not recommended for at least 2 weeks prior to the start of BLINCYTO® treatment, during treatment, and until immune recovery following last cycle of BLINCYTO®.
- **Benzyl Alcohol Toxicity in Neonates:** Serious adverse reactions, including fatal reactions and the "gasping syndrome," have been reported in very low birth weight (VLBW) neonates born weighing less than 1500 g, and early preterm neonates (infants born less than 34 weeks gestational age) who received intravenous drugs containing benzyl alcohol as a preservative. Early preterm VLBW neonates may be more likely to develop these reactions because they may be less able to metabolize benzyl alcohol.

Use the preservative-free preparations of BLINCYTO® where possible in neonates. When prescribing BLINCYTO® (with preservative) for neonatal patients, consider the combined daily metabolic load of benzyl alcohol from all sources including BLINCYTO® (with preservative), other products containing benzyl alcohol or other excipients (e.g., ethanol, propylene glycol) which compete with benzyl alcohol for the same metabolic pathway.

Monitor neonatal patients receiving BLINCYTO® (with preservative) for new or worsening metabolic acidosis. The minimum amount of benzyl alcohol at which serious adverse reactions may occur in neonates is not known. The BLINCYTO® 7-Day bag (with preservative) contains 7.4 mg of benzyl alcohol per mL

Embryo-Fetal Toxicity: Based on its mechanism of action, BLINCYTO® may cause fetal harm when administered to a pregnant woman. Advise pregnant women of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception during treatment with BLINCYTO® and for 48 hours after the last dose.

Adverse Reactions

• The safety of BLINCYTO® in adult and pediatric patients one month and older with MRD-positive B-cell precursor ALL (n=137), relapsed or refractory B-cell precursor ALL (n=267), and Philadelphia chromosome-negative B cell precursor ALL in consolidation (n=165) was evaluated in clinical studies. The most common adverse reactions (≥ 20%) to BLINCYTO® in this pooled population were pyrexia, infusion-related reactions, headache, infection, musculoskeletal pain, neutropenia, nausea, anemia, thrombocytopenia, and diarrhea.

Dosage and Administration Guidelines

- BLINCYTO® is administered as a continuous intravenous infusion at a constant flow rate using an infusion pump which should be programmable, lockable, non-elastomeric, and have an alarm.
- It is very important that the instructions for preparation (including admixing) and administration provided in the full Prescribing Information are strictly followed to minimize medication errors (including underdose and overdose).

Please see accompanying BLINCYTO® full Prescribing Information, including BOXED WARNINGS.

Please note: The information provided in this document is of a general nature and for informational purposes only; it is not intended to be comprehensive or instructive. Coding and coverage policies change periodically and often without warning. The healthcare provider is solely responsible for determining coverage and reimbursement parameters and appropriate coding for their own patients and procedures. In no way should the information provided in this document be considered a guarantee of coverage or reimbursement for any product or service.

SUPPORT SERVICES

AMGEN Support

CALL 866-264-2778

Monday to Friday, 9:00 am to 8:00 pm ET, or visit www.AmgenSupportPlus.com.

We're right here, right when you need us



HCP Support Center

Our Amgen SupportPlus Representatives can assist with issues around patient coverage, prior authorizations, co-pay programs, and more.

Benefits Verification

Verify patient's insurance plan coverage details

Prior Authorization Requirements

· Provide payer-specific prior authorization forms

Amgen SupportPlus Customer Portal

- · A tool for managing patient benefits verification and more
- · Submit, store, and retrieve benefit verifications electronically



Amgen® Patient Navigator

A single point of contact to help answer questions about access and reimbursement, navigating treatment logistics, and to provide supplemental resources as your patients transition from hospital to outpatient care.

Amgen Patient Navigators can help with:

- · Benefits verification and understanding coverage
- Prior authorization process
- · Reimbursement and access resources

The Amgen Patient Navigator is not part of a patient's treatment team and does not provide medical advice or case management services. The Amgen Patient Navigator does not administer Amgen medications. Patients should always consult their healthcare provider regarding medical decisions or treatment concerns.

AMGEN TherapyLocator •

Visit AmgenTherapyLocator.com to locate alternative sites where BLINCYTO® can be administered to your patients*

*The information on this website is reported by independent third-party sites that administer or deliver treatment to patients. It is not comprehensive of all sites that handle the therapies listed, and Amgen does not confirm accuracy or otherwise endorse any of these sites.

Note: Coding and coverage policies change periodically and often without warning. The healthcare provider is solely responsible for determining coverage and reimbursement parameters and appropriate coding for his/her own patients and procedures. This information is not a guarantee of coverage or reimbursement.

AMGEN Support +

1234 5678 9100 0123

MEMBER ID: XXXXXXXXXXXX GROUP: XXXXXXXXXXXX

Questions? Call (866) 264-2778

AMGEN Support Co-Pay Program

The Amgen SupportPlus Co-Pay Program may help eligible patients with private or commercial insurance lower their out-of-pocket costs.

- · Pay as little as \$0 out-of-pocket for each dose or cycle
- Can be applied to deductible, co-insurance, and co-payment[†]
- No income eligibility requirement

†Eligibility criteria and program maximums apply. See AmgenSupportPlus.com/copay for full Terms and Conditions

Encourage your patients with private or commercial insurance to check eligibility and enroll at AmgenSupportPlus.com/copay

What if my patient doesn't have private or commercial insurance?

Amgen SupportPlus can provide your patients with information about independent nonprofit foundations that may be able to help.[‡]

‡Eligibility for resources provided by independent nonprofit patient assistance programs is based on the profit's criteria. Amgen has no control over these programs and provides information as a courtesy only.

References: 1. MLN Matters. Billing for home infusion therapy services on or after January 1, 2021. https://www.cms.gov/files/document/mm11880.pdf. Accessed June 12, 2024. 2. CMS. Medicare Claims Processing Manual Chapter 20. https://www.cms.gov/Regulations-and-Guidance/Guidance/Guidance/Manuals/Downloads/clm104c20.pdf. Accessed June 12, 2024. 3. CMS, CY 2024 Medicare Physician Fee Schedule Rule, Available at: https://www.cms.gov/newsroom/fact-sheets/calendar-year-cy-2024-medicare-physician-fee-schedule-final-rule; Accessed June 3, 2024. 4. CMS, 2% Payment Adjustment Sequestration Changes, December 16, 2021, available at Chapter 20. https://www.cms.gov/Regulations-and-Guidance/Manuals/Downloads/clim104e2/pdf. Accessed June 12, 2024. 3. CMS, SY Payment Agustrant Sequestration Chapters, December 16, 2021, available at https://www.cms.gov/newsroom/facts-heets/celandan-de-schedule-final-rule-accessed June 12, 2024. 5. Value Healthcare Services. Understanding Hospital Revenue Codes. Paths of the Code of the C Advisory Commission. Durable medical equipment payment system. http://www.medpac.gov/docs/default-source/payment-basics/medpac_payment_basics_16_dme_final.pdf. Revised October 2016. Accessed June 3, 2024.

36. CMS. CY 2020 HH PPS Final Rule. https://www.govinfo.gov/content/pkg/FR-2020-11-04/pdf/2020-24146.pdf. Accessed June 3, 2024.

37. Medi-Cal. Provider training 2021: pharmacy billing: home infusion, compound drugs, durable medical equipment & medical supplies. https://files.medi-cal.ca.gov/pubsdoco/outreach_education/workbooks/Workbook_ph-b_dme.pdf. Accessed June 3, 2024.







Billing and Coding Considerations for BLINCYTO®

This Information Sheet is intended to help healthcare professionals understand the key billing and coding considerations for BLINCYTO[®] and its related services and supplies when using the Food and Drug Administration (FDA)-approved dosing options across treatment settings.

Updates regarding Medicare Home Infusion Therapy Benefit:

- **1.** Starting January 1, 2021, Medicare implemented the permanent home infusion therapy benefit that provides separate Part B coverage and payment for qualified home infusion therapy services¹
 - · Medicare updated the codes used to report the provision of home infusion therapy services
 - The new codes differentiate new visits vs subsequent visits for home infusion therapy services
 - Claims for home infusion therapy services will be billed separately from the drug, pump, and other supplies. These services must be reported to the A/B Medicare Administrative Contractor (MAC), and are reimbursed by Medicare at rates set by the Medicare Physician Fee Schedule. Claims for the drug, pump, and supplies should continue being sent to the Durable Medical Equipment (DME) MAC and are payable under the Prosthetics, Orthotics, and Supplies (DMEPOS) Fee Schedule^{1,2}
 - Home infusion therapy services are equal to 5 hours per calendar day, billed in 15 minute increments

Please see pages 10 and 11 for sample claim forms showing coding changes that may be appropriate to report services for Medicare beneficiaries receiving BLINCYTO® treatment via home infusion

2. Due to COVID-19 Public Health Emergency (PHE), Medicare temporarily revised the definition of direct supervision to include the virtual presence of the supervising physician or other qualified healthcare provider using real-time, interactive audio and video telecommunications technology through to December 31, 2024³

Medicare sequestration has been fully reinstated beginning with the third quarter of 2022 and as such, the Medicare portion of payment rates are reduced by 2%.4

Please note that the information in this resource is intended to be educational and is not a guarantee of reimbursement. Coverage, coding, and billing requirements vary by health plan so be sure to check with individual payers for detailed guidance.

INDICATIONS

BLINCYTO® (blinatumomab) is indicated for the treatment of CD19-positive B-cell precursor acute lymphoblastic leukemia (ALL) in adult and pediatric patients one month and older with:

- Philadelphia chromosome-negative disease in the consolidation phase of multiphase chemotherapy
- Minimal residual disease (MRD) greater than or equal to 0.1% in first or second complete remission
- Relapsed or refractory disease

IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGICAL TOXICITIES including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

- Cytokine Release Syndrome (CRS), which may be life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® and treat with corticosteroids as recommended.
- Neurological toxicities, including immune effector cell-associated neurotoxicity syndrome (ICANS) which may be severe, life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® as recommended.

<u>Click here</u> to see BLINCYTO® full Prescribing Information, including **BOXED WARNINGS**. Please see additional Important Safety Information on pages 18-19.



Hospital Inpatient (HIP) Site of Service - Multiple Payers (Medicare and Non-Medicare)

Item	Revenue Code ^{5,6,*}	Coding Information (ICD-10-CM ⁷ /HCPCS ⁸ /CPT ⁹ /ICD-10-PCS ¹⁰)	Notes
Diagnosis: Encounter for drug therapy and ALL	N/A	Z51.12 Encounter for antineoplastic immunotherapy AND C91.00 Acute lymphoblastic leukemia not having achieved remission/failed remission OR C91.01 Acute lymphoblastic leukemia, in remission OR C91.02 Acute lymphoblastic leukemia, in relapse	Report the appropriate ICD-10-CM diagnosis code(s) to describe the patient's condition.
Drug: BLINCYTO® and external infusion pump (EIP)	Report the appropriate revenue code for the cost center in which the service is performed; eg, • Medicare: 0250 General pharmacy • Other payers: 0250 or 0636 Drugs requiring detailed coding (if required by a given payer)	J9039 Injection, blinatumomab, 1 mcg	
	Report the appropriate revenue code for the cost center in which the service is performed; eg, • 0290 DME	E0791 Parenteral infusion pump, stationary, single or multi-channel E0776 IV pole	
Administration: Continuous intravenous infusion (CIVI) via EIP	Report the appropriate revenue code for the cost center in which the service is performed; eg, • 0261 IV therapy: Infusion pump	3E03305 Introduction of other antineoplastic into peripheral vein, percutaneous approach† 0R 3E04305 Introduction of other antineoplastic into central vein, percutaneous approach† 96416 Chemotherapy administration, IV infusion technique; initiation of prolonged chemotherapy infusion (more than 8 hours) requiring use of a portable or implantable pump 0R 96521 Refilling and maintenance of a portable pump	

Coding Information Definitions:

ICD-10-CM – International Classification of Diseases, 10th Revision, Clinical Modification

HCPCS - Healthcare Common Procedure Coding System

CPT - Current Procedural Terminology


ICD-10-PCS – International Classification of Diseases, 10th Revision, Procedure Coding System

^{*}This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.

[†]The previous ICD-10-PCS codes that described the administration of BLINCYTO® (XW03351 and XW04351) have been deleted and should not be used for dates of service on or after October 1, 2021.



Sample UB-04 (CMS-1450) Form: Hospital Inpatient Administration



^{*}This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.

[†]The previous ICD-10-PCS codes that described the administration of BLÍNCYTO® (XW03351 and XW04351) have been deleted and should not be used for dates of service on or after October 1, 2021.



Hospital Outpatient Department (HOPD) - Multiple Payers (Medicare and Non-Medicare)

Item	Revenue Code ^{5,6,*}	Coding Information (ICD-10-CM ⁷ /CPT ⁹ /HCPCS ⁸ /NDC ¹¹)	Notes
Diagnosis: Encounter for drug therapy and ALL	N/A	Z51.12 Encounter for antineoplastic immunotherapy AND C91.00 Acute lymphoblastic leukemia not having achieved remission OR C91.01 Acute lymphoblastic leukemia, in remission OR C91.02 Acute lymphoblastic leukemia, in relapse	Report the appropriate ICD-10-CM diagnosis code(s) to describe the patient's condition.
Procedure: Administration via CIVI using an EIP	Report the appropriate revenue code for the cost center in which the service is performed; eg, • 0261 IV therapy: Infusion pump • 026x IV therapy	96416 Chemotherapy administration, IV infusion technique; initiation of prolonged chemotherapy infusion (more than 8 hours) requiring use of a portable or implantable pump OR 96521 Refilling and maintenance of portable pump OR G0498 Chemotherapy administration, IV infusion technique; initiation of infusion in the office/clinic setting using office/clinic pump/ supplies, with continuation of the infusion in the community setting (eg, home, domiciliary, rest home, or assisted living) using a portable pump provided by the office/clinic; includes follow-up office/clinic visit at the conclusion of the infusion	CPT codes may be used to report the CIVI procedures associated with BLINCYTO® to the Part A/B MAC and non-Medicare payers. For Medicare patients, HCPCS code G0498 will replace CPT and HCPCS codes (96416, E0781, and 99211–99215) previously used to bill for prolonged infusion services when the CIVI is started in the HOPD. It does not apply to BLINCYTO® when the CIVI is started in the inpatient setting or via home infusion.8,9,12 Certain payers may not recognize G0498 and require itemization of specific items, instead. The healthcare provider should consult the payer or MAC to determine which code is most appropriate for administration of BLINCYTO®. If the clinic bills the G-code to the MAC, the cost of the pump and supplies is bundled and should not be billed separately to the DME MAC. ¹³
Drug: BLINCYTO®	Report the appropriate revenue code for the cost center in which the service is performed; eg, • Medicare: 0636 Drug requiring detailed coding • Other payers: 0250 or 0636 General pharmacy (if required by a given payer)	J9039 Injection, blinatumomab, 1 mcg JW Discarded drug/not administered to any patient JZ Zero drug amount discarded/not administered to any patient JG Drug or biological acquired with 340B Drug Pricing Program discount TB Drug or biological acquired with 340B Drug Pricing Program discount	Medicare policies reflect the code for BLINCYTO® (J9039 per 1 mcg) and has a maximum utilization of 210 units per date of service (based on prescribing information).¹⁴ However, coding and coverage requirements may vary by payer. Like many payers, Medicare requires the use of the modifier JW and JZ, which provides payment for the amount of drug or biologic discarded, as well as for the dose administered, up to the amount of the drug or biologic as indicated on the vial or label for a single-dose vial (SDV).¹⁵ Note: Effective for dates of service on or after July 1, 2023, Medicare Part B claims require the use of the new JZ modifier for single-use vials and containers when there are no discarded drug amounts. Medicare claims also continue to require the use of the JW modifier (Drug amount discarded/not administered to any patient) for drugs and biologicals that are separately payable under Medicare Part B with discarded amounts from single-dose containers.¹ Starting January 1, 2024, CMS is requiring all 340B covered entities, including hospital-based and nonhospital-based entities, that submit claims for separately payable Part B drugs and biologicals to report modifier "JG" or "TB" on claim lines for drugs acquired through the 340B Drug Pricing Program. Starting January 1, 2025, 340B covered entities must report the "TB" modifier on claims.¹⁶
	N/A	NDC: 55513016001 BLINCYTO® 35 mcg lyophilized powder, SDV IV solution stabilizer, 10 mL SDV	Some payers (eg, Medicaid) may require listing the NDC in addition to the HCPCS J-code. When reporting the NDC on claims, use the 11-digit NDC in the 5-4-2 format. ¹⁷ Insert a leading zero in the appropriate section to complete the 5-4-2 digit format. Remove the dashes prior to entering the NDC on the claim form.

Coding Information Definition: NDC – National Drug Code

^{*}This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.

[†]Reporting policies for discarded units for payers other than traditional fee-for-service Medicare may vary; providers should check with their specific plans about policies related to billing for discarded drug and use of the JW and JZ modifiers.




Hospital Outpatient Department (HOPD) - Multiple Payers (Medicare and Non-Medicare) (continued)

Item	Revenue Code ^{5,6,*}	Coding Information (HCPCS ⁸)	Notes
DME: EIP and supplies	Report the appropriate revenue code for the cost center in which the service is performed; eg, 0290 DME	E0779 Ambulatory infusion pump, mechanical, reusable, for infusion 8 hours or greater OR E0781 Ambulatory infusion pump, single or multiple channels, electric or battery operated, with administrative equipment, worn by patient OR A4222 Infusion supplies for external drug infusion pump, per cassette or bag Modifiers for use with E-codes for IV pump -KD Drug or biologic infused through DME -RR Rental -KH DMEPOS item, initial claim, purchase or first rental month -KI DMEPOS item, second or third rental months -KJ DMEPOS item, parenteral enteral nutrition (pen) pump or capped rental, fourth to 15th rental months	Please note that Medicare specifically requires DMEPOS accreditation in order to bill a DME MAC. Non-Medicare payers may allow billing for all services and supplies under a medical or other benefit. Report the appropriate EIP code and appropriate modifier(s) as documented in the medical record. Modifiers may be used to provide additional detail when billing for the EIP to the DME MAC. ⁸ Note: Drug administration codes may get billed to the MAC and the E-codes may get billed separately to the DME MAC. Report any supplies as necessary.

^{*}This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.



Sample UB-04 (CMS-1450) Form: Hospital Outpatient Administration



^{*}This is not an all-inclusive list of revenue codes; revenue codes will vary by institution. Revenue codes are only required on the CMS-1450.



Physician Office - Multiple Payers (Medicare and Non-Medicare)

Item	Coding Information (ICD-10-CM ⁷ /CPT ⁹ /HCPCS ⁸ /NDC ¹¹)	Notes
Diagnosis: Encounter for drug therapy and ALL	Z51.12 Encounter for antineoplastic immunotherapy AND C91.00 Acute lymphoblastic leukemia not having achieved remission OR C91.01 Acute lymphoblastic leukemia, in remission OR C91.02 Acute lymphoblastic leukemia, in relapse	Report the appropriate ICD-10-CM code(s) to describe the patient's condition.
Procedure: Administration via CIVI using an EIP	96416 Chemotherapy administration, intravenous infusion technique; initiation of prolonged chemotherapy infusion (more than 8 hours) requiring use of a portable or implantable pump OR 96521 Refilling and maintenance of portable pump OR G0498 Chemotherapy administration, IV infusion technique; initiation of infusion in the office/clinic setting using office/clinic pump/supplies, with continuation of the infusion in the community setting (eg, home, domiciliary, rest home, or assisted living) using a portable pump provided by the office/clinic; includes follow-up office/clinic visit at the conclusion of the infusion	CPT codes may be used to report the CIVI procedures associated with BLINCYTO® to the Part A/B MAC and non-Medicare payers. For Medicare patients, HCPCS code G0498 will replace CPT codes and HCPCS (96416, E0781, and 99211–99215) previously used to bill for prolonged infusion services when the CIVI is started in the physician office. It does not apply to BLINCYTO® when the CIVI is started in the inpatient setting or via home infusion. ^{8,9,12} Some payers may not recognize G0498 and require itemization of specific items, instead. The healthcare provider should consult the payer or MAC to determine which code is most appropriate for administration of BLINCYTO®.
Drug: BLINCYTO®	J9039 Injection, blinatumomab, 1 mcg JW Discarded drug/not administered to any patient JZ Zero drug amount discarded/not administered to any patient	Medicare requires use of the HCPCS code in the physician office setting ¹⁸ and has a maximum utilization of 210 units per date of service (based on prescribing information). However, coding requirements may vary by payer. Like many payers, Medicare requires the use of the modifier JW and JZ, which provides payment for the amount of drug or biologic discarded, as well as for the dose administered, up to the amount of the drug or biologic as indicated on the vial or label for an SDV. Note: Effective for dates of service on or after July 1, 2023, Medicare Part B claims require the use of the new JZ modifier for single-use vials and containers when there are no discarded drug amounts. Medicare claims also continue to require the use of the JW modifier (Drug amount discarded/not administered to any patient) for drugs and biologicals that are separately payable under Medicare Part B with discarded amounts from single-dose containers.*
	NDC: 55513016001 BLINCYTO® 35 mcg lyophilized powder, SDV IV solution stabilizer, 10 mL SDV	Some payers (eg, Medicaid) may require listing the NDC in addition to the HCPCS J-code. When reporting the NDC on claims, use the 11-digit NDC in the 5-4-2 format. ¹⁷ Insert a leading zero in the appropriate section to complete the 5-4-2 digit format. Remove the dashes prior to entering the NDC on the claim form.
DME: EIP and supplies	E0779 Ambulatory infusion pump, mechanical, reusable, for infusion 8 hours or greater ²² E0781 Ambulatory infusion pump, single or multiple channels, electric or battery operated, with administrative equipment, worn by patient ²² G0498 Chemotherapy administration, IV infusion technique; initiation of infusion in the office/clinic setting using office/clinic pump/supplies, with continuation of the infusion in the community setting (eg, home, domiciliary, rest home, or assisted living) using a portable pump provided by the office/clinic; includes follow-up office/clinic visit at the conclusion of the infusion A4222 Infusion supplies for external drug infusion pump, per cassette or bag Modifiers for EIP -KD Drug or biologic infused through DME -RR Rental -KH DMEPOS item, initial claim or first rental month -KI DMEPOS item, second or third rental months -KJ DMEPOS item, fourth to 15th rental months	Report the appropriate EIP code and appropriate modifier(s) as documented in the medical record. Modifiers may be used to provide additional detail when billing for the EIP to the DME MAC. ³ Note: Drug administration codes may get billed to the MAC and the E-codes may get billed separately to the DME MAC. If the office bills the G-code to the MAC, the cost of the pump and supplies is bundled and should not be billed separately to the DME MAC. ¹³ Report any supplies as necessary.

^{*}Reporting policies for discarded units for payers other than traditional fee-for-service Medicare may vary; providers should check with their specific plans about policies related to billing for discarded drug and use of the JW and JZ modifiers.



Sample CMS-1500 Form: Physician Office Administration

	APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NOTICE OF TRANSPORT	UCC) 02/12 CHAMPVA GROUP HEALTH	PLAN — BLK LUNG —	1a. INSURED'S I.D. NUMBE	ا) ٦	PICA For Program in Item 1)
	(Medicaid#) (ID#/DoD#) 2. PATIENT'S NAME (Last Name, First Name, Middle Initial)	(Member ID#) (ID#) 3. PATIENT'S BI	(ID#) (ID#)	4. INSURED'S NAME (Last)	lame, First Name, Mid	dle Initial)
	5. PATIENT'S ADDRESS (No., Street)		M F ATIONSHIP TO INSURED	7. INSURED'S ADDRESS (N	o., Street)	
	СІТУ	STATE 8. RESERVED F	OR NUCC USE	СПУ		STATE
	ZIP CODE TELEPHONE (Include Area	Code)		ZIP CODE	TELEPHONE (Ir	iclude Area Code)
	9. OTHER INSURED'S NAME (Last Name, First Name, Middle	Initial) 10. IS PATIENT'S	S CONDITION RELATED TO:	11. INSURED'S POLICY GR	DUP OR FECA NUME	ER ER
	a. OTHER INSURED'S POLICY OR GROUP NUMBER	a. EMPLOYMEN	T? (Current or Previous)	a. INSURED'S DATE OF BIF	TH Y	SEX F
	b. RESERVED FOR NUCC USE	b. AUTO ACCIDI	PLACE (State)	b. OTHER CLAIM ID (Design		
	c. RESERVED FOR NUCC USE	c. OTHER ACCI	. <mark></mark>	O- F-44b-	RAM NAM	E
	d. INSURANCE PLAN NAME OR PROGRAM NAME	10d. CLAIM COD	piagnosis (Box 2) appropriate diagn ICD-10-CM:		FIT PLAN	ems 9, 9a, and 9d.
	ter the NDC, as UNX (X = the number pecific claim submission requirements 21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Related A. Z51.12 B. C91.02 E. F.	e A-L to service line below (24E	C91.01 Acute ly leukemia, in re C91.02 Acute ly leukemia, in re Final codes deper documentation	mission OR ymphoblastic lapse	ord - En	AGNOSIS POINTER (Box 24E): ter the letter (A–L) that corresponds the diagnosis in Box 21
	1.	D. PROCEDURES, SERVICE (Explain Unusual Circum CPT/HCPCS		F. G	H. I.	UNITS (Box 24G):
1	N455513016001 UNX MM DD YY MM DD YY 11	J9039	A B	XXX XX X	NPI	Report units of service for both units administered and amount of
2	MM DD YY MM DD YY 11	J9039 JW	АВ	xxx xx x	NPI	discarded drug. BLINCYTO® dose reported as 1 unit per mcg. Report
3	MM DD YY MM DD YY 11	96419	A B	XXX XX 1	NPI	1 unit for initiation of CIVI via EIP or refill of EIP
ter the app	RVICE (Box 24B): propriate 2-digit place of that corresponds to the re services are rendered; eg,	Use the CPT code 96416 Chemother	e representing the pro rapy administration, i	ocedure performe intravenous infusi	d, such as ini on technique	ministration procedure. tiation OR refill; eg, ; initiation of prolonged ole or implantable pump



Home Infusion - Multiple Payers (Medicare and Non-Medicare)

Item	Coding Information (ICD-10-CM ⁷ /CPT ⁹ /HCPCS ⁸ /NDC ¹¹)	Notes
Diagnosis: Encounter for drug therapy and ALL	Z51.12 Encounter for antineoplastic immunotherapy AND C91.00 Acute lymphoblastic leukemia not having achieved remission OR C91.01 Acute lymphoblastic leukemia, in remission OR C91.02 Acute lymphoblastic leukemia, in relapse	Report the appropriate ICD-10-CM code(s) to describe the patient's condition.
Procedure: Administration via CIVI using an EIP	G0090 Professional services, initial visit, for the administration of intravenous chemotherapy or other highly complex infusion drug or biological for each infusion drug administration calendar day in the individual's home, each 15 minutes' G0070 Professional services for the administration of intravenous chemotherapy or other intravenous highly complex drug or biological infusion for each infusion drug administration calendar day in the individual's home, each 15 minutes' 99601 Home infusion/specialty drug administration, per visit (up to 2 hours) 99602 Each additional hour S9329 Home infusion therapy, chemotherapy infusion; administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem S9330 Home infusion therapy, continuous (24 hours or more) chemotherapy infusion; administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem S9338 Home infusion therapy, immunotherapy, administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem S9379 Home infusion therapy, infusion therapy, not otherwise classified; administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem	Home infusion therapy services for Medicare beneficiaries receiving BLINCYTO® should be billed using G0090 for an initial visit and G0070 for subsequent visits. Some or all Medicare contractors may reject chemotherapy CPT codes with the availability of G0070 and G0090. These services must be reported to the A/B MAC, and are reimbursed by Medicare at rates set by the Medicare Physician Fee Schedule. They are billed and paid separately from the external infusion pump and drug, which are billed to the DME MAC and reimbursed under the DMEPOS Fee Schedule. Medicare requires that a claim for BLINCYTO® be billed no more than 30 days prior to the visit. Otherwise, payment for the home infusion therapy service will be denied.¹ These services may be covered by Medicaid, commercial plans, or Medicare Advantage plans.²0 CPT codes 99601 and 99602, as well as certain S-codes, may be used to report home infusion therapy services to other payer types other than FFS Medicare. Please note that FFS Medicare does not recognize S-codes, although other payers might.²0
Drug: BLINCYTO®	J9039 Injection, blinatumomab, 1 mcg JW Discarded drug/not administered to any patient JZ Zero drug amount discarded/not administered to any patient	Medicare requires that claims for BLINCYTO®, the pump, and supplies be sent to the DME MACs. Claims for home infusion therapy services must now be submitted separately and are processed by Part A/B MACs.¹ Medicare sets maximum utilization at 875 units of service (UOS), which is equivalent to 25 vials per month in this site of care.²¹ Many payers require the use of the modifier JW and JZ, which provides payment for the amount of drug or biologic discarded, as well as for the dose administered, up to the amount of the drug or biologic as indicated on the vial or label for an SDV.¹⁵ Note: Effective for dates of service on or after July 1, 2023, Medicare Part B claims require the use of the new JZ modifier for single-use vials and containers when there are no discarded drug amounts. Medicare claims also continue to require the use of the JW modifier (Drug amount discarded/not administered to any patient) for drugs and biologicals that are separately payable under Medicare Part B with discarded amounts from single-dose containers.*
	NDC: 55513016001 BLINCYTO® 35 mcg lyophilized powder, SDV IV solution stabilizer, 10 mL SDV	Some payers (eg, Medicaid) may require listing the NDC in addition to the HCPCS J-code. When reporting the NDC on claims, use the 11-digit NDC in the 5-4-2 format. ¹⁷ Insert a leading zero in the appropriate section to complete the 5-4-2 digit format. Remove the dashes prior to entering the NDC on the claim form.
DME: EIP and supplies	E0779 Ambulatory infusion pump, mechanical, reusable, for infusion 8 hours or greater ²² E0781 Ambulatory infusion pump, single or multiple channels, electric or battery operated, with administrative equipment, worn by patient ²² A4222 Infusion supplies for external drug infusion pump, per cassette or bag Modifiers for EIP -KD Drug or biologic infused through DME -RR Rental -KH DMEPOS item, initial claim or first rental month -KI DMEPOS item, second or third rental months -KJ DMEPOS item, fourth to 15th rental months	Report the appropriate EIP code and appropriate modifier(s) as documented in the medical record. Modifiers may be used to provide additional detail when billing for the EIP to the DME MAC.8 Report any supplies as necessary.

^{*}Reporting policies for discarded units for payers other than traditional fee-for-service Medicare may vary; providers should check with their specific plans about policies related to billing for discarded drug and use of the JW and JZ modifiers.



Sample CMS-1500 Form: Medicare DME MAC for BLINCYTO®, Pump, and Related Supplies by DME Supplier

HEALTH INSURANCE CLAIM FOR APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NU		PICA [← CAPRIER →		
MEDICARE MEDICAID TRICARE (Madicare#) (Madicaid#) (ID#/DoD#) 2. PATIENT'S NAME (Last Name, First Name, Middle Initial)	CHAMPVA GROUP EECA OTHER (Member (D#)) (D#) (D#) (D#) (D#) (D#) (D#) (D#	Insured's I.D. NUMBER (For Program in Item 1) 4. INSURED'S NAME (Last Name, First Name, Middle Initial)			
sample claim form on page 11 for guidance on	6. PATIENT RELATIONSHIP TO INSURED Self Spouse Child Other STATE 8. RESERVED FOR NUCC USE initial) 10. IS PATIENT'S CONDITION RELATED TO: a. EMPLOYMENT? (Current or Previous)	7. INSURED'S ADDRESS (No., Street) CITY STATE ZIP CODE TELEPHONE (Include Area Code) () 11. INSURED'S POLICY GROUP OR FECA NUMBER	RED INFORMATION		
b. RESERVED FOR NUCC USE	b. AUTO ACCIDENT? PLACE (State)	a. INSURED'S DATE OF BIRTH SEX MM DD YY M F b. OTHER CLAIM ID (Designated by NUCC)	AND INSURED		
c. RESERVED FOR NUCC USE	c. OTHER ACCIDENT?	c. INSURANCE PLAN NAME OR PROGRAM NAME			
the 11-digit format. The unit of measure can also be reported, 3 spaces after the NDC, as UNX (X = the number of vials)	INPLETING & SIGNING THIS FORM. IMPLETING & SIGNING THIS FORM. Ithorize the release of any medical or other information necessary selfus either to myself or to the party who accepts assignment DATE	d. IS THERE ANOTHER HEALTH BENEFIT PLAN? YES NO If yes, complete items 9, 9a, and 9d. 13. INSURED'S OR AUTHORIZED PERSON'S SIGNATURE Lauthorize payment of medical benefits to the undersigned physician or supplier for services described below. SIGNED	PATIENT **		
11 DATE OF CURRENT ILLNESS, INJURY, or PREGNANCY (I	MP) 15. OTHER DATE MM DD YY	16. DATES PATIENT LINABLE TO WORK IN CURRENT OCCUPATION	<u></u>		
1. NAME OF REFERRING PROVIDER OR OTHER SOURCE 1. ADDITIONAL CLAIM INFORMATION (Designated by NUCC) 2. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate 2. L	C91.00 Acute lymphoblastic leuk	stic immunotherapy AND kemia not having achieved remission OR kemia, in remission OR kemia, in relapse			
From To PLACE OF	D. PROCEDURES, SERVICES, OR SUPPLIES (Explain Unusual Circumstances) CPT/HCPCS I MODIFIER POINTER	F. G. H. I. PSTT ID UNITS (Box 240 S) CHARGES UNITS (Box 240 D) CONTROL OF THE PROPERTY WINTS OF THE PROPERTY			
1 N455513016001 UNX		units administ	f service for both ered and amount		
MM DD YY MM DD YY 12 18 19 19 19 19 19 19 19	J9039 A B	XXX XX X NPI of discarded d	rug. BLINCYTO® as 1 unit per mcg.		
MM DD YY MM DD YY 12	J9039 JW A B	XXX XX X NPI Report 1 unit e	ach for EIP and		
MM DD YY MM DD YY 12	E0781 RR KH A B	XXX XX 1 NPI other supplies			
4 MM DD YY MM DD YY 12	A4222 A B	DIAGNOSIS POINTER (Box 24E): Enter the le			
5 1 1 1		that corresponds to the diagnosis in box 2			
PLACE OF SERVICE (Box 24B): Enter the appropriate 2-digit place of service code that corresponds to the location where services are rendered; eg, 12 Home	Drug: J9039 for BLINCYTO* JW/JZ Discard Modifier: JW or JZ mo DME external infusion pump claims single-use containers	odes and modifiers; eg, difier required for Medicare including infused drugs in	C Use		
9	 IV Pump: E0781 Ambulatory infus A4222 Infusion supplies for exterior 				
	NUCC Instruction Manual available at: www.nucc. Other codes may be appropriate. Check with individual Medicare DME MACs for detailed guidance ORM 1500 (02-12)				



Sample CMS-1500 form: Medicare A/B MAC for Home Infusion Therapy Services by Home Infusion Therapy Supplier

HEALTH INSURANCE CLAIM FORM APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 02/12			CARRIER →
1. MEDICARE MEDICAID TRICARE CHAMPV (Medicare#) (Medicaid#) (//D#/D-D#) (Member II	— HEALTH PLAN — BLK LUNG —	1a. INSURED'S I.D. NUMBER (F	For Program In Item 1)
In 2021, Medicare requires separate claims for home infusion therapy services and for drugs furnished as items of DME in the home infusion setting. This sample claim shows an example for billing home infusion therapy services for a Medicare patient. See the sample claim form on page 10 for guidance on billing for drugs furnished as an item of DME for a Medicare beneficiary	3. PATIENT'S BIRTH DATE SEX M F G. PATIENT RELATIONSHIP TO INSURED Self Spouse Child Other 8. RESERVED FOR NUCC USE	INSURED'S NAME (Last Name, First Name, Mid INSURED'S ADDRESS (No., Street) CITY ZIP CODE TELEPHONE (Ir. () 11. INSURED'S POLICY GROUP OR FECA NUMB	STATE OLUMB Area Code) ER VILLE STATE
a. OTHER INSURED'S POLICY OR GROUP NUMBER	a. EMPLOYMENT? (Current or Previous) YES NO	a. INSURED'S DATE OF BIRTH	SEX PER PROPERTY OF THE PROPER
b. RESERVED FOR NUCC USE	b. AUTO ACCIDENT? PLACE (State)	b. OTHER CLAIM ID (Designated by NUCC)	AND IN
c. RESERVED FOR NUCC USE	c. OTHER ACCIDENT?	c. INSURANCE PLAN NAME OR PROGRAM NAM	
d. INSURANCE PLAN NAME OR PROGRAM NAME	10d. CLAIM CODES (Designated by NUCC)	d. IS THERE ANOTHER HEALTH BENEFIT PLANS YES NO If yes, complete its	
READ BACK OF FORM BEFORE COMPLETIN. 12. PATIENT'S OR AUTHORIZED PERSON'S SIGNATURE I authorize the to process this claim. I also request payment of government benefits either below. SIGNED	release of any medical or other information necessary	 INSURED'S OR AUTHORIZED PERSON'S SIG payment of medical benefits to the undersigned services described below. SIGNED	
14. DATE OF CURRENT ILLNESS, INJURY, or PREGNANCY (LMP) DD YOUAL. 17. NAME OF REFERRING PROVIDER OR OTHER SOURCE 19. ADDITIONAL CLAIM INFORMATION (Designated by NUCC) 21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate A-L to s A. Z51.12 B. C91.02 E. L. F. G	DIAGNOSIS (BOX 21): Enter the appropriate diagnosis Z51.12 Encounter for antineoplas C91.00 Acute lymphoblastic leul C91.01 Acute lymphoblastic leul C91.02 Acute lymphoblastic leul Final codes depend on medical	stic immunotherapy AND kemia not having achieved remi kemia, in remission OR kemia, in relapse	ssion OR
I	L. L. E. E. E.	F. G. H. I. DAYS PSUT	HAUTO (Day 040)
1 MM DD YY MM DD YY SERVICE EMG CPT/HCP		\$ CHARGES UNITS Pen QUAL.	UNITS (Box 24G): Report units of service for the administration of
2		XXX XX X NPI	BLINCYTO [®] , reported as 1 unit per 15 minutes of
3 MM DD YY MM DD YY 12 J9039	A B	0 00 X NPI	time of IV infusion
PLACE OF SERVICE (Box 24B): Enter the appropriate 2-digit place of service code that corresponds to the location where services are rendered; eq.	DIAGNOSIS POINTER letter (A–L) that co diagnosis in Box 21	(Box 24E): Enter the rresponds to the	PHYSICIAN OR SU
12 Home 26. PAT 27. SIGNATURE OF PHYSICIAN OR SUPPLIER INCLUDING DEGREES OR CREDENTIALS (I certify that the statements on the reverse apply to this bill and are made a part thereof.) 28. SEF	OCEDURES/SERVICES/SUPPLIES (E ter the appropriate CPT/HCPCS of IV Infusion: G0090 for IV infusion Drug: J9039 is added to identify List a zero charge to indicate the her codes may be appropriate. C	codes and modifiers; eg, n of BLINCYTO®, initial visit BLINCYTO® as the drug related at no reimbursement for the dru	to the administration service; g is expected



Sample CMS-1500 Form: Non-Medicare Payer by Home Infusion Provider

	HEALTH INSURANCE CLAIM FORM APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) (27)	2				
	PICA PICA					
	1. MEDICARE MEDICAID TRICARE CHAMM (Medicare#) (Medicaid#) (ID#/DoD#) (Member	rID#) HEALTH PLAN BLK LUNG (ID#) (ID#)		tem 1)		
	2. PATIENT'S NAME (Last Name, First Name, Middle Initial)	3. PATIENT'S BIRTH DATE SEX MM DD YY M F	4. INSURED'S NAME (Last Name, First Name, Middle Initial)			
	5. PATIENT'S ADDRESS (No., Street)	6. PATIENT RELATIONSHIP TO INSURED Self Spouse Child Other	7. INSURED'S ADDRESS (No., Street)			
	CITY STAT		CITY ST	TATE		
	ZIP CODE TELEPHONE (Include Area Code)		ZIP CODE TELEPHONE (Include Area Cod	de)		
	9. OTHER INSURED'S NAME (Last Name, First Name, Middle Initial)	10. IS PATIENT'S CONDITION RELATED TO:	11, INSURED'S POLICY GROUP OR FECA NUMBER			
	a. OTHER INSURED'S POLICY OR GROUP NUMBER	a. EMPLOYMENT? (Current or Previous)	a. INSURED'S DATE OF BIRTH SEX			
	b. RESERVED FOR NUCC USE	b. AUTO ACCIDENT? NO PLACE (State)	b. OTHER CLAIM ID (Designated by NUCC)			
	c. RESERVED FOR NUCC USE	YES NO OTHER ACCIDENT?	c. INSURANCE PLAN NAME OR PROGRAM NAME			
	d. INSURANCE PLAN NAME OR PROGRAM NAME	YES NO				
		10d. CLAIM CODES (Designated by NUCC)	d. IS THERE ANOTHER HEALTH BENEFIT PLAN? YES NO If yes, complete items 9, 9a, and			
	12. PATIENT'S OR AUTHORIZED PERSON'S SIGNATURE I authorize it to noncess this claim. Lalso request payment of government benefits eith	e release of any medical or other information necessary	 INSURED'S OR AUTHORIZED PERSON'S SIGNATURE I auth payment of medical benefits to the undersigned physician or su services described below. 			
	A SHADED AREA): When required by the payer, DC qualifier "N4," indicating that an NDC	_ DATE	SIGNED			
follows, and	the NDC in the 11-digit format. The unit of	HER DATE MM DD YY	16. DATES PATIENT UNABLE TO WORK IN CURRENT OCCUPA	ITION		
UNX (X = th	n also be reported, 3 spaces after the NDC, as e number of vials). Verify the payer-specific		nter the appropriate diagnosis code; eg, l r antineoplastic immunotherapy AND	CD-10-CM:		
claim submi	ssion requirements	C91.00 Acute lymph	oblastic leukemia not having achieved re	emission OR		
	21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate A-L to se	rvice line belov • C91.02 Acute lymph	noblastic leukemia, in remission OR noblastic leukemia, in relapse			
	A. Z51.12 B. C91.02 C.	Final codes depend or	n medical record documentation			
	I J K.		F. G. H. I. DAYS			
		olain Unusual Circumstances) PCS MODIFIER POIL TER		R (Box 24E): Enter the		
	MM DD YY MM DD YY 12 J903	9 A B	xxx xx x diagnosis in Box 2			
	MM DD YY MM DD YY 12 J903	9 JW AB	XXX XX X IIIITS (Per 24C)			
	MM DD YY MM DD YY 12 996	01 AB		service for both units		
	4 MM DD YY MM DD YY 12 A42	22 A B	of drug adminis	tered and amount ug. BLINCYTO® dose		
	IVIVI DD 11 IVIVI DD 11 12 A42	AD	reported with 1	unit per 1 mcg		
Enter the ap	orresponds to the location where rendered; eg,	nd modifiers; eg, Drug: J9039 for BLINCYTO® 99601 Home infusion/specialty of A4222 Infusion supplies for exter option ther codes may be appropriate. Ch	ox 24D): Enter the appropriate CPT/HCPC Irug administration, per visit (up to 2 hour and drug infusion pump, per cassette or ineck with individual payers for detailed good and the control of the contr	irs) infusion uidance		
	Se	rvice Medicare may vary; provide	rs should check with their specific plans ded drug and use of the JW and JZ modif	about		



BLINCYTO® Dosing Options¹¹

Dosing option	Dose per vial X number of SDVs*	Number of billing units
24-hour	35 mcg X 1 vial	35
48-hour	35 mcg X 1-2 vials	35-70
7-day	35 mcg X 4-6 vials	140-210

^{*}Number of SDVs depends on dose, infusion duration, and patient's weight.11

Key Considerations for the BLINCYTO® 7-day Infusion Option (7-DIO)



Minor variations are expected in coding, billing, and claims filing for the BLINCYTO® 7-DIO.²⁰



The 7-DIO requires 6 vials of BLINCYTO® and 1 vial of IV Solution Stabilizer for patients ≥ 45 kg. For patients weighing less than 45 kg, 4 to 5 vials are required. The safety of the administration of BLINCYTO® at a BSA of less than 0.4 m² has not been established.¹¹ Refer to the Prescribing Information for details on handling and preparation.



If the units field on a claim form cannot accommodate more than 99 units, utilize multiple lines to capture all units (eg, 99+98+13). Payers may require separate reporting of drug units administered and discarded.²⁰



Less frequent claim submissions are expected with utilization of the 7-DIO. Typically the entire 7-DIO can be billed on the day of administration/refill. However, be sure to refer to payer guidelines for maximum daily quantity limits. Apply the appropriate dates of service as needed.²⁰



If the 7-DIO is interrupted mid-treatment, refer to payer guidelines for reporting and documentation in these cases. If full reimbursement is withheld by the payer, refer to Amgen's Product Return Policy for assistance.



Existing codes and modifiers are adequate to report BLINCYTO® and its related services; however, payer requirements may vary with respect to:²⁰

- The entities that can bill for DME and the associated supplies
- The number of units billed for BLINCYTO® J9039 (HCPCS units vs number of vials)
- Covered diagnosis codes
- Covered nursing services (eg, infusion services at patient's home)
- Drug claim submission options (eg, 1 or more dates of service on claims)
- Reporting policies for discarded units for payers other than traditional fee-for-service Medicare
 may vary; providers should check with their specific plans about policies related to billing for
 discarded drug and use of the JW and JZ modifiers.

UNDERSTANDING EXAMPLES OF



REIMBURSEMENT ACROSS SITES OF CARE

A BLINCYTO® patient transitions through multiple sites of care. This guide shows how major payers in the United States (commercial plans, Medicare, and Medicaid) offer coverage in each setting and reimburse for each component of care:



Drug



Pump and Supplies



Hospitalization



Professional Services (ie, drug administration)

INDICATIONS

BLINCYTO® (blinatumomab) is indicated for the treatment of CD19-positive B-cell precursor acute lymphoblastic leukemia (ALL) in adult and pediatric patients one month and older with:

- Philadelphia chromosome-negative disease in the consolidation phase of multiphase chemotherapy
- Minimal residual disease (MRD) greater than or equal to 0.1% in first or second complete remission
- Relapsed or refractory disease


IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGICAL TOXICITIES including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

- Cytokine Release Syndrome (CRS), which may be life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® and treat with corticosteroids as recommended.
- Neurological toxicities, including immune effector cell-associated neurotoxicity syndrome (ICANS) which
 may be severe, life-threatening or fatal, occurred in patients receiving BLINCYTO[®]. Interrupt or discontinue
 BLINCYTO[®] as recommended.

BLINCYTO® (blinatumomab) Reimbursement Process

Coverage of BLINCYTO® and its administration is required in all these sites of care to avoid interruption in treatment.



The scenarios depicted above illustrate the most common ones for accessing BLINCYTO® via the buy-and-bill acquisition process, where the entity that acquires the product also administers it to the patient.

BLINCYTO® can also be acquired via a specialty pharmacy provider, including:

- Third-party specialty pharmacies that contract with a payer to supply specialty products covered under the medical benefit
- Specialty pharmacies owned by hospitals, physician offices, ambulatory infusion clinics, and/or home infusion companies that may also administer the medication

BLINCYTO® Reimbursement Across Transitions in Site of Care

BLINCYTO®-eligible patients need coverage for the following: Drug + Pump + Hospitalization + Administration

Inpatient Hospital					
Components of BLINCYTO® Care	Commercially Insured Patients	FFS Medicare Patients	FFS Medicaid Patients		
BLINCYTO®	MS-DRG-based or global payment; typically includes BLINCYTO ^{®23}	MS-DRG payment includes BLINCYTO® ²⁴	APR-DRG-based payment; typically includes BLINCYTO®27		
Ē	Reimbursement varies by contracts between providers and payers	Covered under Medicare Part A benefit ²⁵	Reimbursement varies by state; may follow Medicare		
Pump and Supplies	Some hospitals, in their contracts with managed care organizations, may negotiate a "carve out" benefit for drugs such as BLINCYTO® • May allow separate payment of such drugs outside of the bundled payment for inpatient services	Hospital may be eligible for outlier payments if cost of admission exceeds certain threshold	allowable amounts		
Hospitalization		Reimbursement varies for the 11 IPPS-Exempt Cancer Hospitals ²⁶			
Professional Services	Physician services may be covered separately outside of the bundled payment	Physician services may be covered and reimbursed according to the MPFS under Medicare Part B benefit	Physician services may be covered and paid outside of the bundled payment		

Key: APR-DRG-All Patient-Refined Diagnosis Related Groups; FFS-fee-for-service; IPPS-Inpatient Prospective Payment System; MPFS-Medicare Physician Fee Schedule; MS-DRG-Medicare Severity Diagnosis-Related Group.

Outpatient Hospital					
Components of BLINCYTO® Care	Commercially Insured Patients	FFS Medicare Patients	FFS Medicaid Patients		
BLINCYTO®	Reimbursed based on contracted rates; methodology varies Examples: ²⁸ • ASP + X% • WAC + X% • AWP-X% May need prior authorization	Covered under Medicare Part B benefit Typically reimbursed based on ASP + 6% when administered in a 340B hospital setting (with 2% sequestration reduction) ^{29,30} MUE cap of 210 units (approx. 6 vials) per date of service applies ^{31,8}	Reimbursement may be similar to Medicare OR State-defined limit (eg, California uses a federal upper limit) ³² May need prior authorization		
Pump and Supplies	Reimbursement is bundled into the payment for the infusion service	Covered under Medicare Part B benefit Reimbursement is bundled into the payment for the infusion service	Reimbursed based on fee schedule or bundled into the payment for the infusion service Rates vary by state		
Professional Services	Reimbursed based on contracted rate	Reimbursed based on the Medicare OPPS			

 $Key: ASP-average \ sales \ price; AWP-average \ wholesale \ price; FFS-fee-for-service; \ MUE-medically \ unlikely \ edit; OPPS-Outpatient \ Prospective \ Payment \ System; \ WAC-wholesale \ acquisition \ cost.$

Note: The information here describes coverage and payment for BLINCYTO® under FFS Medicare and FFS Medicaid. Coverage and payment for patients enrolled in Medicare Advantage and/or Medicaid managed care organizations varies widely and is often similar to commercial insurance.

Physician Office					
Components of BLINCYTO® Care	Commercially Insured Patients	FFS Medicare Patients	FFS Medicaid Patients		
BLINCYTO®	Reimbursed based on contracted rates; methodology varies Examples: ²⁸ • ASP + X% • WAC + X% • AWP – X% May need prior authorization	Covered under Medicare Part B benefit Typically reimbursed based on ASP + 6% (with 2% sequestration reduction) ^{30,33} MUE cap of 210 units (approx. 6 vials) per date of service applies ^{8,19}	Reimbursement may be similar to Medicare OR State-defined limit (eg, California uses a federal upper limit) 32 May need prior authorization		
Pump and Supplies	Reimbursed based on contracted rate and bundled into payment for the infusion service	Covered under Medicare Part B benefit Reimbursement is bundled into the payment for the infusion service	Typically reimbursed based on fee schedule or bundled into the payment for the infusion service Rates vary by state		
Professional Services	Reimbursed based on contracted rate	Reimbursed based on the MPFS			

Key: ASP-average sales price; AWP-average wholesale price; FFS-fee-for-service; MPFS-Medicare Physician Fee Schedule; MUE-medically unlikely edit; WAC-wholesale acquisition cost.

Home Infusion			
Components of BLINCYTO® Care	Commercially Insured Patients	FFS Medicare Patients	FFS Medicaid Patients
BLINCYTO®	Reimbursed based on contracted rates; methodology varies Examples: ²⁸ • ASP + X% • WAC + X% • AWP-X% May need prior authorization	Covered under Medicare Part B as long as it is supplied in a covered external infusion pump and the IV is initiated in home infusion setting ³⁴ Typically reimbursed based on ASP + 6% (with 2% sequestration reduction) ^{30,33} Billing cap of 25 vials per month applies ²¹	Reimbursement may be similar to Medicare OR State-defined limit (eg, California uses a federal upper limit) ³⁷ May need prior authorization
Pump and Supplies	Reimbursed based on contracted rate	Covered under Medicare Part B benefit Reimbursed as part of the Medicare DMEPOS Fee Schedule ³⁵	Typically reimbursed based on fee schedule Rates vary by state
Professional Services	Reimbursed based on contracted rate	Covered under Part B Reimbursed under the home infusion therapy services benefit in 15-minute increments for applicable providers ³⁶	

Key: ASP-average sales price; AWP-average wholesale price; DMEPOS-Durable Medical Equipment Prosthetics, Orthotics, and Supplies; FFS-fee-for-service; WAC-wholesale acquisition cost.

 ${\sf Note: Medicare\ home\ infusion\ benefit\ is\ distinct\ and\ separate\ from\ the\ Medicare\ home\ health\ benefit.}$



BLINCYTO® Indications and Important Safety Information



INDICATIONS

BLINCYTO® (blinatumomab) is indicated for the treatment of CD19-positive B-cell precursor acute lymphoblastic leukemia (ALL) in adult and pediatric patients one month and older with:

- Philadelphia chromosome-negative disease in the consolidation phase of multiphase chemotherapy
- · Minimal residual disease (MRD) greater than or equal to 0.1% in first or second complete remission
- Relapsed or refractory disease

IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGICAL TOXICITIES including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

- Cytokine Release Syndrome (CRS), which may be life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® and treat with corticosteroids as recommended.
- Neurological toxicities, including immune effector cell-associated neurotoxicity syndrome (ICANS) which may be severe, life-threatening or fatal, occurred in patients receiving BLINCYTO®. Interrupt or discontinue BLINCYTO® as recommended.

Contraindications

BLINCYTO® is contraindicated in patients with a known hypersensitivity to blinatumomab or to any component of the product formulation.

Warnings and Precautions

- Cytokine Release Syndrome (CRS): CRS, which may be life-threatening or fatal, occurred in patients receiving BLINCYTO®. The median time to onset of CRS is 2 days after the start of infusion and the median time to resolution of CRS was 5 days among cases that resolved. Closely monitor and advise patients to contact their healthcare professional for signs and symptoms of serious adverse events such as fever, headache, nausea, asthenia, hypotension, increased alanine aminotransferase (ALT), increased aspartate aminotransferase (AST), increased total bilirubin, and disseminated intravascular coagulation (DIC). The manifestations of CRS after treatment with BLINCYTO® overlap with those of infusion reactions, capillary leak syndrome (CLS), and hemophagocytic histiocytosis/macrophage activation syndrome (MAS). Using all of these terms to define CRS in clinical trials of BLINCYTO, CRS was reported in 15% of patients with R/R ALL, in 7% of patients with MRD-positive ALL, and in 16% of patients receiving BLINCYTO® cycles in the consolidation phase of therapy. If severe CRS occurs, interrupt BLINCYTO® until CRS resolves. Discontinue BLINCYTO® permanently if life-threatening CRS occurs. Administer corticosteroids for severe or life-threatening CRS.
- Neurological Toxicities, including Immune Effector Cell-Associated Neurotoxicity Syndrome: BLINCYTO® can cause serious or life-threatening neurologic toxicity, including ICANS. The incidence of neurologic toxicities in clinical trials was approximately 65%. The median time to the first event was within the first 2 weeks of BLINCYTO® treatment. The most common (≥ 10%) manifestations of neurological toxicity were headache and tremor. Grade 3 or higher neurological toxicities occurred in approximately 13% of patients, including encephalopathy, convulsions, speech disorders, disturbances in consciousness, confusion and disorientation, and coordination and balance disorders. Manifestations of neurological toxicity included cranial nerve disorders. The majority of neurologic toxicities resolved following interruption of BLINCYTO®, but some resulted in treatment discontinuation.
 - The incidence of signs and symptoms consistent with ICANS in clinical trials was 7.5%. The onset of ICANS can be concurrent with CRS, following resolution of CRS, or in the absence of CRS. There is limited experience with BLINCYTO® in patients with active ALL in the central nervous system (CNS) or a history of neurologic events. Patients with a history or presence of clinically relevant CNS pathology were excluded from clinical studies. Patients with Down Syndrome over the age of 10 years may have a higher risk of seizures with BLINCYTO® therapy.
 - Monitor patients for signs and symptoms of neurological toxicities, including ICANS, and interrupt or discontinue BLINCYTO® as outlined in the PI. Advise outpatients to contact their healthcare professional if they develop signs or symptoms of neurological toxicities.
- Infections: Approximately 25% of patients receiving BLINCYTO® in clinical trials experienced serious infections such as sepsis, pneumonia, bacteremia, opportunistic infections, and catheter-site infections, some of which were life-threatening or fatal. Administer prophylactic antibiotics and employ surveillance testing as appropriate during treatment. Monitor patients for signs or symptoms of infection and treat appropriately, including interruption or discontinuation of BLINCYTO® as needed.
- Tumor Lysis Syndrome (TLS), which may be life-threatening or fatal, has been observed. Preventive measures, including pretreatment nontoxic cytoreduction and on-treatment hydration, should be used during BLINCYTO® treatment. Monitor patients for signs and symptoms of TLS and interrupt or discontinue BLINCYTO® as needed to manage these events.
- Neutropenia and Febrile Neutropenia, including life-threatening cases, have been observed. Monitor appropriate laboratory parameters
 (including, but not limited to, white blood cell count and absolute neutrophil count) during BLINCYTO® infusion and interrupt BLINCYTO® if
 prolonged neutropenia occurs.
- Effects on Ability to Drive and Use Machines: Due to the possibility of neurological events, including seizures and ICANS, patients receiving BLINCYTO® are at risk for loss of consciousness, and should be advised against driving and engaging in hazardous occupations or activities such as operating heavy or potentially dangerous machinery while BLINCYTO® is being administered.
- Elevated Liver Enzymes: Transient elevations in liver enzymes have been associated with BLINCYTO® treatment with a median time to onset of 3 days. In patients receiving BLINCYTO®, although the majority of these events were observed in the setting of CRS, some cases of elevated liver enzymes were observed outside the setting of CRS, with a median time to onset of 19 days. Grade 3 or greater elevations in liver enzymes occurred in approximately 7% of patients outside the setting of CRS and resulted in treatment discontinuation in less than



IMPORTANT SAFETY INFORMATION (continued)

1% of patients. Monitor ALT, AST, gamma-glutamyl transferase, and total blood bilirubin prior to the start of and during BLINCYTO® treatment. BLINCYTO® treatment should be interrupted if transaminases rise to > 5 times the upper limit of normal (ULN) or if total bilirubin rises to > 3 times ULN.

- Pancreatitis: Fatal pancreatitis has been reported in patients receiving BLINCYTO® in combination with dexamethasone in clinical trials and the post-marketing setting. Evaluate patients who develop signs and symptoms of pancreatitis and interrupt or discontinue BLINCYTO® and dexamethasone as needed.
- **Leukoencephalopathy:** Although the clinical significance is unknown, cranial magnetic resonance imaging (MRI) changes showing leukoencephalopathy have been observed in patients receiving BLINCYTO®, especially in patients previously treated with cranial irradiation and antileukemic chemotherapy.
- **Preparation and administration** errors have occurred with BLINCYTO® treatment. Follow instructions for preparation (including admixing) and administration in the PI strictly to minimize medication errors (including underdose and overdose).
- Immunization: Vaccination with live virus vaccines is not recommended for at least 2 weeks prior to the start of BLINCYTO® treatment, during treatment, and until immune recovery following last cycle of BLINCYTO®.
- Benzyl Alcohol Toxicity in Neonates: Serious adverse reactions, including fatal reactions and the "gasping syndrome," have been reported in very low birth weight (VLBW) neonates born weighing less than 1500 g, and early preterm neonates (infants born less than 34 weeks gestational age) who received intravenous drugs containing benzyl alcohol as a preservative. Early preterm VLBW neonates may be more likely to develop these reactions because they may be less able to metabolize benzyl alcohol.
 - Use the preservative-free preparations of BLINCYTO® where possible in neonates. When prescribing BLINCYTO® (with preservative) for neonatal patients, consider the combined daily metabolic load of benzyl alcohol from all sources including BLINCYTO® (with preservative), other products containing benzyl alcohol or other excipients (e.g., ethanol, propylene glycol) which compete with benzyl alcohol for the same metabolic pathway.
 - Monitor neonatal patients receiving BLINCYTO® (with preservative) for new or worsening metabolic acidosis. The minimum amount of benzyl alcohol at which serious adverse reactions may occur in neonates is not known. The BLINCYTO® 7-Day bag (with preservative) contains 7.4 mg of benzyl alcohol per mL
- **Embryo-Fetal Toxicity:** Based on its mechanism of action, BLINCYTO® may cause fetal harm when administered to a pregnant woman. Advise pregnant women of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception during treatment with BLINCYTO® and for 48 hours after the last dose.

Adverse Reactions

• The safety of BLINCYTO® in adult and pediatric patients one month and older with MRD-positive B-cell precursor ALL (n=137), relapsed or refractory B-cell precursor ALL (n=267), and Philadelphia chromosome-negative B cell precursor ALL in consolidation (n=165) was evaluated in clinical studies. The most common adverse reactions (≥ 20%) to BLINCYTO® in this pooled population were pyrexia, infusion-related reactions, headache, infection, musculoskeletal pain, neutropenia, nausea, anemia, thrombocytopenia, and diarrhea.

Dosage and Administration Guidelines

- BLINCYTO® is administered as a continuous intravenous infusion at a constant flow rate using an infusion pump which should be programmable, lockable, non-elastomeric, and have an alarm.
- It is very important that the instructions for preparation (including admixing) and administration provided in the full Prescribing Information are strictly followed to minimize medication errors (including underdose and overdose).

Please see BLINCYTO® full Prescribing Information, including BOXED WARNINGS.

Please note: The information provided in this document is of a general nature and for informational purposes only; it is not intended to be comprehensive or instructive. Coding and coverage policies change periodically and often without warning. The healthcare provider is solely responsible for determining coverage and reimbursement parameters and appropriate coding for their own patients and procedures. In no way should the information provided in this document be considered a guarantee of coverage or reimbursement for any product or service.

SUPPORT SERVICES



CALL 866-264-2778

Monday to Friday, 9:00 am to 8:00 pm ET, or visit www.AmgenSupportPlus.com.

We're right here, right when you need us



HCP Support Center

Our Amgen SupportPlus Representatives can assist with issues around patient coverage, prior authorizations, co-pay programs, and more.

Benefits Verification

Verify patient's insurance plan coverage details

Prior Authorization Requirements

· Provide payer-specific prior authorization forms

Amgen SupportPlus Customer Portal

- · A tool for managing patient benefits verification and more
- Submit, store, and retrieve benefit verifications electronically



Amgen® Patient Navigator

A single point of contact to help answer questions about access and reimbursement, navigating treatment logistics, and to provide supplemental resources as your patients transition from hospital to outpatient care.

Amgen Patient Navigators can help with:

- · Benefits verification and understanding coverage
- · Prior authorization process
- · Reimbursement and access resources

The Amgen Patient Navigator is not part of a patient's treatment team and does not provide medical advice or case management services. The Amgen Patient Navigator does not administer Amgen medications. Patients should always consult their healthcare provider regarding medical decisions or treatment concerns.

AMGEN TherapyLocator •

Visit **AmgenTherapyLocator.com** to locate alternative sites where BLINCYTO® can be administered to your patients*

*The information on this website is reported by independent third-party sites that administer or deliver treatment to patients. It is not comprehensive of all sites that handle the therapies listed, and Amgen does not confirm accuracy or otherwise endorse any of these sites.

Note: Coding and coverage policies change periodically and often without warning. The healthcare provider is solely responsible for determining coverage and reimbursement parameters and appropriate coding for his/her own patients and procedures. This information is not a guarantee of coverage or reimbursement.

AMGEN Support⁺

1234 5678 9100 0123

RxBIN: XXXXXX

MEMBER ID: XXXXXXXXXXX GOOD 00/1
GROUP: XXXXXXXXXXX

Questions? Call (866) 264-2778

AMGEN Support | Co-Pay Program

The Amgen SupportPlus Co-Pay Program may help eligible patients with private or commercial insurance lower their out-of-pocket costs.

- · Pay as little as \$0 out-of-pocket for each dose or cycle
- Can be applied to deductible, co-insurance, and co-payment[†]
- No income eligibility requirement

†Eligibility criteria and program maximums apply. See AmgenSupportPlus.com/copay for full Terms and Conditions

Encourage your patients with private or commercial insurance to check eligibility and enroll at AmgenSupportPlus.com/copay

What if my patient doesn't have private or commercial insurance?

Amgen SupportPlus can provide your patients with information about independent nonprofit foundations that may be able to help. ‡

[‡]Eligibility for resources provided by independent nonprofit patient assistance programs is based on the nonprofit's criteria. Amgen has no control over these programs and provides information as a courtesy only.

References: 1.MLM Matters. Billing for home infusion therapy services on or after January 1, 2021. https://www.cms.gov/flees/document/mmt180.pdf. Accessed June 12, 2024. A. CMS, CY 2024. Medicare Psysician Fee Schedule Brit. Psychiatry 1, 2024. https://www.cms.gov/flees/document/mmt180.pdf. Accessed June 12, 2024. A. CMS, CY 2024. Medicare Psysician Fee Schedule Brit. Psychiatry 1, 2024. https://www.cms.gov/flees/document/mmt180.pdf. Accessed June 12, 2024. A. CMS, CY 2024. Medicare Psysician Fee Schedule Brit. Psychiatry 1, 2024. https://www.cms.gov/flees/document/mmt180.pdf. Accessed June 12, 2024. https://www.cms.gov/pub/Helahls.psychule/he



